Contents lists available at ScienceDirect



European Journal of Operational Research

journal homepage: www.elsevier.com/locate/ejor



CrossMark

Continuous Optimization

Continuous multifacility ordered median location problems *

Víctor Blanco^{a,*}, Justo Puerto^b, Safae El-Haj Ben-Ali^c

^a Department of Quantitative Methods for Economics & Business, Universidad de Granada, 18011 Granada, Spain
 ^b Department de Estadística e Investigación Operativa, Universidad de Sevilla, 41012 Sevilla, Spain
 ^c Département Génie Electrique et Informatique, USMBA University, ENSA-Fès, Morocco

ARTICLE INFO

Article history: Received 24 October 2014 Accepted 28 October 2015 Available online 9 November 2015

Keywords: Continuous multifacility location Ordered median problems Semidefinite programming Second order cone programming

ABSTRACT

In this paper we propose a general methodology for solving a broad class of continuous, multifacility location problems, in any dimension and with ℓ_{τ} -norms proposing two different methodologies: (1) by a new second order cone mixed integer programming formulation and (2) by formulating a sequence of semidefinite programs that converges to the solution of the problem; each of these relaxed problems solvable with SDP solvers in polynomial time.

© 2015 Elsevier B.V. and Association of European Operational Research Societies (EURO) within the International Federation of Operational Research Societies (IFORS). All rights reserved.

1. Introduction

Multifacility location problems are among the most interesting and difficult problems in Location Analysis. It is well-known that even in their discrete version the *p*-median and *p*-center problems are already NP-hard (see Kariv and Hakimi, 1979.) A lot of attention has been paid in the last decades to these classes of problems, namely location-allocation problems, since they are easy to describe and to understand and they still capture the essence of difficult problems in combinatorial optimization. A comprehensive overview over existing models and their applications is given by Drezner and Hamacher (2002) and the references therein.

On the other hand, also in the last two decades locators have devoted much effort to solve continuous location problems that fall within the general class of global optimization, i.e. convexity properties are lost. Given a set of demand points (existing facilities) the goal is to locate several facilities to provide service to the existing ones (demand points) minimizing some globalizing function of the travel distances. Assuming that each demand point will be served by its closest facility we are faced with another location-allocation problem but now the new facilities can be located anywhere in the

* Corresponding author. Tel.: +34 958249637.

framework space and therefore they are not confined to be in an "a priori" given set of locations. These problems are much harder than the discrete ones and not much has been obtained regarding algorithms, and general convergence results, although some exceptions can be found in the literature (see Albareda, Hinojosa, and Puerto, 2015; Brimberg, Drezner, Mladenović, and Salhi, 2014; Brimberg, Hansen, Mladenovic, and Taillard, 2000; Michelot, 1987 and the references therein).

Since the 1990s a new family of objective functions has started to be considered in the area of Location Analysis: the ordered median problem (Nickel & Puerto, 2005). Ordered median problems represent as special cases nearly all classical objective functions in location theory, including the Median, CentDian, Center and k-Centra. Hence, handling the most important objective functions in Location Analysis is possible with one unique model and also new ones may be created by adapting the parameters adequately. More precisely, the p-facility ordered median problem can be formulated as follows: a vector of weights $(\lambda_1, \ldots, \lambda_n)$ is given. The problem is to find locations for the facilities that minimize the weighted sum of distances to the facilities where the distance to the closest point to its allocated facility is multiplied by the weight λ_n , the distance to the second closest, by λ_{n-1} , and so on. The distance to the farthest point is multiplied by λ_1 . As mentioned above, many location problems can be formulated as the ordered 1-median problem by selecting appropriate weights. For example, the vector for which all $\lambda_i = 1$ is the *p*median problem, the problem where $\lambda_1 = 1$ and all others are equal to zero is the *p*-center problem, the problem where $\lambda_1 = \ldots = \lambda_k = 1$ and all others are equal to zero is the *p*-*k*-centrum. Minimizing the range of distances is achieved by $\lambda_1 = 1$, $\lambda_n = -1$ and all others are zero. Lots of results have been obtained for these problems in

http://dx.doi.org/10.1016/j.ejor.2015.10.065

0377-2217/© 2015 Elsevier B.V. and Association of European Operational Research Societies (EURO) within the International Federation of Operational Research Societies (IFORS). All rights reserved.

^{*} The first and second authors were partially supported by the project FQM-5849 (Junta de Andalucía\FEDER). The authors were partially supported by the project MTM2010-19576-C02-01 (MICINN, Spain). The second author also would like to acknowledge networking support by the COST Action TD1207. The authors would like to thank the anonymous referees for their useful comments and suggestions that helped to improve this work.

E-mail addresses: vblanco@ugr.es (V. Blanco), puerto@us.es (J. Puerto), anasafae@gmail.com (S.E.-H. Ben-Ali).

discrete settings, on networks and even in the continuous single facility case (see the book Nickel and Puerto, 2005 and the recent papers Blanco, El-Haj Ben-Ali, and Puerto, 2013; Boland, Dominguez-Marin, Nickel, and Puerto, 2006; Kalcsics, Nickel, Puerto, and Rodríguez-Chía, 2010; Marín, Nickel, Puerto, and Velten, 2009; Puerto, D. Pérez-Brito, and Garcia-González, 2014; Puerto and Rodriguez-Chia, 2011). However, very little is known in the continuous multifacility counterpart (Kalcsics, Nickel, Pozo, Puerto, & Rodriguez-Chia, 2014).

In this paper, we address the multifacility continuous ordered median problem in finite dimension *d* and for general ℓ_{τ} -norm for measuring the distances between points. We show how these problems can be cast within a general family of polynomial optimization problems. Then, we show how these problems can be formulated as mixed integer second order cone programs or, using tools borrowed from the Theory of Moments (Lasserre, 2009b), they can be solved (approximated up to any degree of accuracy) by a series of relaxed problems, each one of them is a SDP, polynomially solvable. We report some computational results on a battery of problems solved with the presented models.

2. Preliminaries

In this section we recall the main definitions and results on Semidefinite Programming and the Theory of Moments that will be useful for the development through this paper. We use standard notation in those fields (see e.g. Lasserre, 2009b; Wolkowicz, Saigal, & Vandenberghe, 2000).

Semidefinite programming (SDP) is relatively a new subfield of convex optimization and probably one of the most exciting development in mathematical programming. SDP is a particular case of conic programming when one considers the convex cone of positive semidefinite matrices, whereas linear programming considers the positive orthant, a polyhedral convex cone. This class of problems theoretically includes a large number of convex programming models such as convex quadratic programming (QP), or second-order cone programming (SOCP). Moreover, semidefinite programming has experienced a great growth during the 1990s, as a consequence of the adaptation of some polynomial time interior point methods firstly developed for linear optimization to solve SDP. SDP duality is not always strong because of the nonlinear positive semidefinite constraint. To avoid duality gaps, we can require the problem and its dual to satisfy some constraint gualification. The purpose of a constraint qualification is to ensure the existence of Lagrange multipliers at optimality in nonlinear problems. These multipliers are an optimal solution for the dual problem, and thus the constraint qualification ensures that strong duality holds: it is possible to achieve primal and dual feasibility with no duality gap. One common choice of constraint gualification is Slater's constraint gualification (Boyd & Vanderberghe, 2004). The handbooks (Anjos & Lasserre, 2012) and (Wolkowicz et al., 2000) provide excellent coverage of SDP as well as extensive bibliographic reviews of the literature up to the year 2012.

On the other hand, the Theory of Moments is a powerful technique that can be applied to approximate hard global optimization problems (Lasserre, 2009b). It can be used to construct hierarchies of approximate problems whose optimal values converge to the optimal value of the multifacility location problems considered in this paper. For this reason next we describe its basic elements. We denote by $\mathbb{R}[x]$ the ring of real polynomials in the variables $x = (x_1, \ldots, x_d)$, for $d \in \mathbb{N}$ ($d \ge 1$), and by $\mathbb{R}[x]_r \subset \mathbb{R}[x]$ the vector space of polynomials of degree at most $r \in \mathbb{N}$ (here \mathbb{N} denotes the set of non-negative integers). We also denote by $B = \{x^{\alpha} : \alpha \in \mathbb{N}^d\}$ a canonical basis of monomials for $\mathbb{R}[x]$, where $x^{\alpha} = x_1^{\alpha_1} \cdots x_d^{\alpha_d}$, for any $\alpha \in \mathbb{N}^d$. Note that $B_r = \{x^{\alpha} \in B : \sum_{i=1}^d \alpha_i \le r\}$ is a basis for $\mathbb{R}[x]_r$. For any sequence indexed in the canonical monomial basis B, $\mathbf{y} = (y_{\alpha})_{\alpha \in \mathbb{N}^d} \subset \mathbb{R}$, let $L_{\mathbf{y}} : \mathbb{R}[x] \to \mathbb{R}$ be the linear functional defined, for any $f = \sum_{\alpha \in \mathbb{N}^d} f_{\alpha} x^{\alpha} \in \mathbb{R}[x]$, as $L_{\mathbf{y}}(f) := \sum_{\alpha \in \mathbb{N}^d} f_{\alpha} y_{\alpha}$.

The *moment* matrix $M_r(\mathbf{y})$ of order r associated with \mathbf{y} , has its rows and columns indexed by the elements in the basis $B = \{x^{\alpha} : \alpha \in \mathbb{N}^d\}$ and for two elements in such a basis, $b_1 = x^{\alpha}, b_2 = x^{\beta}, M_r(\mathbf{y})(b_1, b_2) = M_r(\mathbf{y})(\alpha, \beta) := L_{\mathbf{y}}(x^{\alpha+\beta}) = y_{\alpha+\beta}$, for $|\alpha|, |\beta| \le r$ (here |a| stands for the sum of the coordinates of $a \in \mathbb{N}^d$). Note that the moment matrix of order r has dimension $\binom{d+r}{d} \times \binom{d+r}{d}$ and that there are $\binom{d+2r}{d} \mathbf{y}_{\alpha}$ variables.

there are $\binom{d+2r}{d} \mathbf{y}_{\alpha}$ variables. For $g \in \mathbb{R}[x]$ ($= \sum_{\gamma \in \mathbb{N}^d} g_{\gamma} x^{\gamma}$), the *localizing* matrix $M_r(g\mathbf{y})$ of order r associated with \mathbf{y} and g, has its rows and columns indexed by the elements in B and for $b_1 = x^{\alpha}$, $b_2 = x^{\beta}$, $M_r(g\mathbf{y})(b_1, b_2) = M_r(g\mathbf{y})(\alpha, \beta) := L_{\mathbf{y}}(x^{\alpha+\beta}g(x)) = \sum_{\gamma} g_{\gamma} y_{\gamma+\alpha+\beta}$, for $|\alpha|, |\beta| \le r$.

Observe that a different choice for the basis of $\mathbb{R}[x]$, instead of the standard monomial basis, would give different moment and localizing matrices useful for instance to derive improvements in the relaxations for structured (symmetric or sparse) problems.

The main assumption to be imposed when one wants to assure convergence of some SDP relaxations for solving polynomial optimization problems is known as the Arquimedean property (see for instance Lasserre, 2009a; 2009b) and it is a consequence of Putinar's results (Putinar, 1993).

The importance of Archimedean property stems from the link between such a condition with the semidefiniteness of the moment and localizing matrices (see Putinar, 1993). The use of this property for the particular problems that we deal with through this paper will be given in the next sections. A detailed presentation and an account of its implications can be found by Lasserre (2001). In the following we shall use the following convention: for any matrix $A \in \mathbb{R}^{n \times n}$, we denote by $A \succeq 0$ (resp. $A \succ 0$) that the matrix A is positive semidefinite (resp. positive definite).

Proposition 1 Lasserre (2001). Let $\mathbf{K} := \{x \in \mathbb{R}^d : g_j(x) \ge 0, j = 1, ..., m\} \subset \mathbb{R}^d$ satisfy the Archimedean Property and let $p \in \mathbb{R}[X]$ be a polynomial. Let $r \ge r_0 := \max\{\lceil \deg \frac{p}{2} \rceil, \lceil \deg \frac{g_1}{2} \rceil, ..., \lceil \deg \frac{g_m}{2} \rceil\}$, and consider the hierarchy of semidefinite relaxations

$$Q_r: \begin{array}{ccc} \inf_{y} \mathbf{L}_{\mathbf{y}}(p) \\ Q_r: & \mathbf{M}_r(\mathbf{y}) \succeq 0, \\ \mathbf{M}_{r-\lceil \deg g_j/2 \rceil}(g_j \mathbf{y}) \succeq 0, & 1 \le j \le m \\ \mathbf{L}_{\mathbf{y}}(y_0) = 1 \end{array}$$
(1)

with optimal value denoted by infQ_r.

Then, the hierarchy of SDP-relaxations $\{(Q_r)_{r \ge r_0}\}$ is monotone nondecreasing and converges to $\rho^* := \min_{x \in K} p(x)$.

SOCP is a particular class of problems that falls within the family of SDP. SOCP refers to convex optimization problems in which a linear function is minimized over the intersection of an affine linear manifold with the Cartesian product of second-order cones. Many applications in engineering, statistics, robust optimization, finance or combinatorial optimization use SOCP models (see Lobo, Vandenberghe, Boyd, & Lebret, 1998). Although the available algorithms for SDP are clearly applicable to SOCP, specialized approaches for SOCP have been proved to be more efficient for this family of problems (see Alizadeh & Goldfarb, 2003).

For the sake of readability and to do the paper self contained we recall the following result, proved by Blanco, Puerto, and El-Haj Ben-Ali (2014b), that allows to represent several types of constraints as intersection of second order cone constraints.

Lemma 2 Blanco et al. (2014b). Let $r, s \in \mathbb{N} \setminus \{0\}$ such that gcd (r, s) = 1. Then, the constraints

$$u^{s}t^{r-s} \ge x^{r},$$

 $x, u, t \ge 0,$

can be equivalently written as a set of at most $\lfloor \log_2(r) \rfloor$ second order cone inequalities with $\lfloor \log_2(r) \rfloor$ new (auxiliary) nonnegative variables.

3. The multiple allocation multifacility ordered median location problem

This section deals with multifacility location models where more than one new facility have to be located to improve the service for the demand points. Several results obtained in previous papers are extended or reformulated with great generality giving a panorama view of the geometric insights of location theory.

We start by considering a class of multifacility ordered median problems already introduced by Nickel and Puerto (2005) and Rodríguez-Chía, Nickel, Puerto, and Fernández (2000). We shall extend these models, originally considered only in dimension 2 and with polyhedral norms to the more general case of any dimension d > 1 and any ℓ_{τ} -norm being $\tau \in \mathbb{Q}, \tau \ge 1$ (here ℓ_{τ} stands for the norm $||x||_{\tau} = (\sum_{i=1}^{d} |x_i|^{\tau})^{\frac{1}{\tau}}$, for all $x \in \mathbb{R}^d$). Unlike the original approaches for the planar case by Carrizosa and Puerto (1995); Nickel and Puerto (2005); Rodríguez-Chía et al. (2000), where even for polyhedral norms there are proposed iterative algorithms for which polynomiality results can not be proven; we shall follow on a different approach. In this section, we provide efficient reformulations of these classes of multifacility continuous location models. We apply tools borrowed from conic programming to prove that these problems can be polynomially solved.

We are given a set of demand points $\{a_1, \ldots, a_n\}$ and three sets of scalars $\{\omega_1, \ldots, \omega_n\} \subseteq \mathbb{R}_+$, $\{\lambda_{11}, \ldots, \lambda_{np}\} \subseteq \mathbb{R}_+$ and $\{\mu_{12}, \mu_{13}, \ldots, \mu_{p-1p}\} \subseteq \mathbb{R}_+$.

The elements ω_i are weights corresponding to the importance given to the existing facilities $a_i, i \in \{1, ..., n\}$ and depending on the choice of the λ -weights we get different classes of problems. The μ -weights represent the penalty per distance unit given when locating two different facilities. We denote by P_n the set of permutations of the first *n* natural numbers (i.e., $P_n = \{\sigma : \{1, ..., n\} \rightarrow \{1, ..., n\}$: σ bijective}).

Natural extensions of the multifacility models considered by Nickel and Puerto (2005) and Rodríguez-Chía et al. (2000), assume that one is looking for the location of *p* new facilities rather than only one. In this formulation the new facilities are chosen to provide service to all the existing facilities minimizing an ordered median objective function. These ordered problems are of course harder to handle than the classical ones not considering ordered distances. To simplify the presentation we consider that the different demand points use the same norm to measure distances, although all our results extend further to the case of mixed norms (different norm for each facility).

Let us consider a set of demand points $\{a_1, a_2, ..., a_n\} \subset \mathbb{R}^d$. We want to locate p new facilities $X = \{x_1, x_2, ..., x_p\}$ which minimize the following expression:

$$f_{\lambda}^{NI}(x_1, x_2, \dots, x_p) = \sum_{i=1}^n \sum_{j=1}^p \lambda_{ij} d_{(i)}(x_j) + \sum_{j=1}^{p-1} \sum_{j'=j+1}^p \mu_{jj'} \|x_j - x_{j'}\|_{\tau},$$
(2)

where for any $x \in \mathbb{R}^d$, $d_i(x) = \omega_i || a_i - x ||_{\tau}$ and $d_{(i)}(x)$ is the *i*th element in the permutation of $(d_1(x), \ldots, d_n(x))$ such that $d_{(1)}(x) \ge d_{(2)}(x) \ge \ldots \ge d_{(i)}(x) \ge \ldots \ge d_{(n)}(x)$. In this model, it is assumed that (see Rodríguez-Chía et al., 2000)

$$\lambda_{1j} \ge \lambda_{2j} \ge \ldots \ge \lambda_{nj} \ge 0, \ \forall j = 1, \ldots, p.$$
 (3)

 $\mu_{jj'} \ge 0$ for any j, j' = 1, ..., p and, as mention above, $d_{(i)}(x_j)$ is the expression, which appears at the *i*th position in the ordered version of the list

$$L_j^{NI} := (w_1 \| x_j - a_1 \|_{\tau}, \dots, w_n \| x_j - a_n \|_{\tau}) \quad \text{for } j = 1, 2, \dots, p.$$
(4)

Note that in this formulation we assign the lambda parameters with respect to each new facility, i.e., x_j is considered to be non-interchangeable with x_i whenever $i \neq j$. For this reason we say that this model has non-interchangeable facilities.

The problem consists of

$$\rho_{\lambda}^{NI} := \min_{x} \{ f_{\lambda}^{NI}(x) : x = (x_1, \dots, x_p), \ x_j \in \mathbb{R}^d, \ \forall j = 1, \dots, p \},$$
(LOCOMF-NI)

The reader should observe that this is the extension of Problem (3) by Rodríguez-Chía et al. (2000, Section 4.1).

Theorem 3. The problem LOCOMF-NI attains an optimal solution.

Proof. We know that

$$\sum_{i=1}^{n} \sum_{j=1}^{p} \lambda_{ij} d_{(i)}(x_j) = \sum_{i=1}^{n} \max_{\sigma} \sum_{j=1}^{p} \lambda_{ij} w_{\sigma(i)} \| x_j - a_{\sigma(i)} \|_{\tau}$$

where σ is a permutation of the set $\{1, 2, ..., p\}$. Therefore, the first part of the objective function is a sum of maxima of convex functions. Hence, it is a convex function. Thus, f_{λ}^{NI} is a convex function as a sum of convex functions.

Next, suppose that we restrict ourselves to consider the problem where $x_j = x$ for all j = 1, ..., p. Assume that x^* is a solution. Then, for any x not optimal it must exist $i_x \in \{1, ..., n\}$ such that $||x - a_{i_x}||_{\tau} \ge ||x^* - a_{i_x}||_{\tau}$. Thus taking x = 0, we get $||x^* - a_{i_0}||_{\tau} \le ||a_{i_0}||_{\tau}$ which implies that $||x^*||_{\tau} - \max_{i=1,...,n} ||a_i||_{\tau} \le$ $\max_{i=1,...,n} ||a_i||_{\tau}$. Hence, any candidate x^* to optimal solution must satisfy $||x^*||_{\tau} \le 2 \max_{i=1,...,n} ||a_i||_{\tau} = M$.

Let us denote by X the set $\{x \in \mathbb{R}^{p \times d} : ||x_j||_{\tau} \le M, \forall j = 1, ..., p\}$. Observe that for solving LOCOMF-NI we can restrict, without loss of generality, to X. Thus, it consists of minimizing the convex function f_{λ}^{NI} over X which is compact in $\mathbb{R}^{p \times d}$ and consequently by Weierstrass theorem, Problem LOCOMF-NI admits at least one optimal solution.

The following result states a sufficient condition for uniqueness of optimal solution of Problem (LOCOMF-NI).

Theorem 4. Assume that $\tau \notin \{1, +\infty\}$, the demand points in A are not collinear and for all i = 1, ..., n there exists at least one $j \in \{1, ..., p\}$ such that $\lambda_{ij} \neq 0$. Then the optimal solution of Problem (LOCOMF-NI) is unique.

Proof. We prove that the function $f_{\lambda}^{NI}(x_1, x_2, ..., x_p)$ is strictly convex which clearly implies the result.

Since the function $f_{\lambda}^{NI}(x_1, x_2, ..., x_p)$ is globally convex we can take without loss of generality two points \bar{x} , \hat{x} and $[\bar{x}, \hat{x}]$ in the same ordered region and $\gamma \in (0, 1)$. (Observe that otherwise, we can decompose the segment in subsegments satisfying the hypothesis and we apply the same argument in each of them.) In the ordered region the sorting of the distances is fixed and to simplify the presentation we suppose that this sorting is induced by the identity permutation.

By hypothesis, the demand points in *A* are not collinear and the $\|\cdot\|_{\tau}$ -norm is strictly convex for $\tau \in (1, +\infty)$; thus, for each $j \in 1, ..., p$ there exists i(j) such that $\|\gamma \bar{x}_j + (1-\gamma)\hat{x}_j - a_{i(j)}\|_{\tau} < \gamma \|\bar{x}_j - a_{i(j)}\|_{\tau} + (1-\gamma)\|\hat{x}_j - a_{i(j)}\|_{\tau}$, $j \in 1, ..., p$. These inequalities imply that $f_{\lambda}^{NI}(\gamma(\bar{x}_1, \bar{x}_2, ..., \bar{x}_p) + (1-\gamma)(\hat{x}_1, \hat{x}_2, ..., \hat{x}_p)) < \gamma f_{\lambda}^{NI}(\bar{x}_1, \bar{x}_2, ..., \bar{x}_p) + (1-\gamma)f_{\lambda}^{NI}(\hat{x}_1, \hat{x}_2, ..., \hat{x}_p)$.

Note that when the demand points in *A* are collinear one can find multiple solutions to Problem (LOCOMF-NI). The following example illustrates this situation.

Example 5. Let $A = \{(0, 0), (1, 1), (2, 2), (10, 10)\}$ be a set of demand points in \mathbb{R}^2 endowed with the ℓ_2 -norm and consider the following weights $w_i = 1$ for all i = 1, ..., 4, $\mu_{12} = 0$ and

$$\lambda_{11} = \lambda_{21} = \lambda_{31} = \lambda_{41} = 1,$$

 $\lambda_{12} = 1, \ \lambda_{22} = \lambda_{32} = \lambda_{42} = 0.$



Fig. 1. Demand points in Example 5 (circles) and solutions (triangles and square).

The optimal value of (LOCOMF-NI) for p = 2 facilities is $f^* = 16\sqrt{2}$ which is attained by $x_1^* \in \{(x, x) : x \in [1, 2]\}, x_2^* = (5, 5)$. Fig. 1 shows the demand points (circles) and the optimal solutions to the problem (triangles and square).

Next we prove that Problem LOCOMF-NI can be reformulated in an appropriate way that allows us the development of an efficient algorithm based on the theory of second order cone programming.

Theorem 6. Let $\tau = \frac{r}{s}$ be such that $r, s \in \mathbb{N} \setminus \{0\}, r \ge s$ and gcd (r, s) = 1. For any set of lambda weights satisfying $\lambda_{1j} \ge ... \ge \lambda_{nj} \ge 0$ for all j = 1, ..., p, Problem (LOCOMF-NI) is equivalent to

$$\rho_{\lambda}^{NI} = \min \sum_{i=1}^{n} \sum_{j=1}^{p} \nu_{ij} + \sum_{\ell=1}^{n} \sum_{j=1}^{p} w_{\ell j} + \sum_{j=1}^{p-1} \sum_{j'=j+1}^{p} t_{jj'}$$
(NIMFOMP) _{λ}

s.t.
$$v_{ij} + w_{\ell j} \ge \lambda_{\ell j} u_{ij}, \forall i, \ell = 1, \dots, n, j = 1, \dots, p$$
 (5)

$$y_{ijk} - x_{jk} + a_{ik} \ge 0, \forall i = 1, ..., n, \ j = 1, ..., p, \ k = 1, ..., d$$
 (6)

$$y_{ijk} + x_{jk} - a_{ik} \ge 0, \forall i = 1, ..., n, \quad j = 1, ..., p, \ k = 1, ..., d,$$
 (7)

$$y_{ijk}^r \le \zeta_{ijk}^s u_{ij}^{r-s}, \forall i = 1, ..., n, \quad j = 1, ..., p, \ k = 1, ..., d,$$
 (8)

$$\omega_i^{\frac{r}{s}} \sum_{k=1}^d \varsigma_{ijk} \le u_{ij}, \forall i = 1, \dots, n, \ j = 1, \dots, p$$
(9)

$$z_{jj'k} - x_{jk} + x_{j'k} \ge 0, \forall j, j' = 1, \dots, p, \ k = 1, \dots, d,$$
(10)

$$z_{jj'k} + x_{jk} - x_{j'k} \ge 0, \forall j, j' = 1, \dots, p, \ k = 1, \dots, d,$$
(11)

$$z_{jj'k}^r \le \xi_{jj'k}^s t_{jj'}^{r-s}, \forall j, j' = 1, \dots, p, \ k = 1, \dots, d,$$
(12)

$$\mu_{jj'}^{\frac{r}{s}} \sum_{k=1}^{d} \xi_{jj'k} \le t_{jj'}, \forall j, j' = 1, \dots, p,$$
(13)

$$\zeta_{ijk} \ge 0, \forall i = 1, ..., n, \ j = 1, ..., p, \ k = 1, ..., d,$$
(14)

$$\xi_{jj'k} \ge 0, \forall j, j' = 1, \dots, p, \ k = 1, \dots, d,$$
 (15)

$$v_{ij} \in \mathbb{R}, w_{\ell j} \in \mathbb{R}, t_{jj'} \ge 0, \forall i = 1, \dots, n, j, j' = 1, \dots, p$$
(16)

$$y_{ijk} \ge 0, z_{jj'k} \ge 0, \forall i = 1, ..., n, j, j' = 1, ..., p, k = 1, ..., d.$$
 (17)

Moreover, $(\text{NIMFOMP}_{\lambda})$ satisfies Slater condition and it can be represented as a second order cone program with $(np + p^2)(2d + 1) + p^2$ linear inequalities and at most $4(p^2d + npd) \log r$ second order cone inequalities.

Proof. Note that the condition $\lambda_{1j} \ge ... \ge \lambda_{nj}$ for all j = 1, ..., p, allows us to write Problem (LOCOMF-NI) as

$$\min_{\mathbf{x}\in\mathbb{R}^{d_p}}\max_{\sigma\in\mathcal{P}_n} \sum_{i=1}^n \sum_{j=1}^p \lambda_{ij}\omega_{\sigma(i)} \|\mathbf{x}_j - a_{\sigma(i)}\|_{\tau} + \sum_{j=1}^{p-1} \sum_{j'=j+1}^p \mu_{jj'} \|\mathbf{x}_j - \mathbf{x}_{j'}\|_{\tau},$$
(18)

Let us introduce the auxiliary variables u_{ij} and $t_{jj'}$, i = 1, ..., n and j, j' = 1, ..., p to which we impose that $u_{ij} \ge \omega_i ||x_j - a_i||_{\tau}$ and $t_{jj'} \ge \mu_{jj'} ||x_j - x_{j'}||_{\tau}$, to model the problem in a convenient form.

Now, for any permutation $\sigma \in P_n$, let $u_{\sigma j} = (u_{\sigma(1)j}, \ldots, u_{\sigma(n)j})$ for $j = 1, \ldots, p$. Moreover, let us denote by (·) the permutation that sorts any vector in nonincreasing sequence, i.e. $u_{(1)j} \ge u_{(2)j} \ge \ldots \ge u_{(n)j}$. Using that $\lambda_{1j} \ge \ldots \ge \lambda_{nj}$ and since $u_{ij} \ge 0$, for all $i = 1, \ldots, n$ and $j = 1, \ldots, p$ then

$$\sum_{i=1}^{n} \sum_{j=1}^{p} \lambda_{ij} u_{(i)j} = \max_{\sigma \in \mathcal{P}_n} \sum_{i=1}^{n} \sum_{j=1}^{p} \lambda_{ij} u_{\sigma(i)j}.$$
(19)

The permutations in P_n can be represented by the binary variables

$$p_{ijk} = \begin{cases} 1, & \text{if } u_{ij} \text{ goes in position } k, \\ 0, & \text{otherwise,} \end{cases}$$

imposing that they verify the following constraints:

$$\sum_{i=1}^{n} p_{ijk} = 1, \ \forall j = 1, \dots, p, \ k = 1, \dots, n,$$
$$\sum_{k=1}^{n} p_{ijk} = 1, \ \forall i = 1, \dots, n, \ j = 1, \dots, p.$$

Next, combining the two sets of variables we obtain that righthand side of (19) can be equivalently written as

$$\begin{cases} \sum_{i=1}^{n} \sum_{j=1}^{p} \lambda_{ij} u_{(i)j} = \max \sum_{i=1}^{n} \sum_{j=1}^{p} \sum_{k=1}^{n} \lambda_{ij} u_{ij} p_{ijk} \\ s.t \sum_{i=1}^{n} p_{ijk} = 1, \ \forall j = 1, \dots, p, \ k = 1, \dots, n, \\ \sum_{k=1}^{n} p_{ijk} = 1, \ \forall i = 1, \dots, n, \ j = 1, \dots, p, \\ p_{ijk} \in \{0, 1\}. \end{cases}$$

$$(20)$$

Now, we point out that for fixed *j*, i.e. considering the terms that contain u_{1j}, \ldots, u_{nj} , we have that

$$\begin{cases} \sum_{i=1}^{n} \lambda_{ij} u_{(i)j} = \max \sum_{i=1}^{n} \sum_{k=1}^{n} \lambda_{ij} u_{ij} p_{ijk} \\ s.t \sum_{i=1}^{n} p_{ijk} = 1, \ \forall k = 1, \dots, n, \\ \sum_{i=1}^{n} p_{ijk} = 1, \ \forall i = 1, \dots, n, \\ p_{ijk} \in \{0, 1\}. \end{cases}$$
(21)

The problem above is an assignment problem and its constraint matrix is totally unimodular, so that solving a continuous relaxation of the problem always yields an integral solution vector (see Ahuja, Magnanti, & Orlin, 1993), and thus a valid permutation. Moreover, the dual of the linear programming relaxation of (21) is strong and also gives the value of the original binary formulation of (21). Hence, for fixed $j \in \{1, ..., p\}$ and for any vector $u_{.j} \in \mathbb{R}^n$, by using the dual of

the assignment problem (21) we obtain the following expression

$$\begin{cases} \sum_{i=1}^{n} \lambda_{ij} u_{(i)j} = \min \sum_{i=1}^{n} v_{ij} + \sum_{l=1}^{n} w_{lj} \\ s.t \quad v_{ij} + w_{lj} \ge \lambda_{lj} u_{ij}, \ \forall i, l = 1, \dots, n. \end{cases}$$
(22)

Finally, we replace (22) in (18) and we get

$$\begin{cases} \min\sum_{i=1}^{n}\sum_{j=1}^{p}v_{ij} + \sum_{l=1}^{n}\sum_{j=1}^{p}w_{lj} + \sum_{j=1}^{p-1}\sum_{j'=j+1}^{p}t_{jj'} \\ \text{s.t.} \quad v_{ij} + w_{lj} \ge \lambda_{lj}u_{ij}, \\ u_{ij} \ge \omega_{i}||\mathbf{x}_{j} - q_{i}||_{\tau}, \\ t_{jj'} \ge \mu_{jj'}||\mathbf{x}_{j} - \mathbf{x}_{j'}||_{\tau}, \end{cases} \quad \forall i, l = 1, \dots, n, \ j = 1, \dots, p, \\ i = 1, \dots, n, \ j = 1, \dots, p, \\ j, j' = 1, \dots, p. \end{cases}$$

$$(23)$$

It remains to prove that each inequality $u_{ij} \ge \omega_i ||x_j - a_i||_{\tau}$, i = 1, ..., n, j = 1, ..., p can be replaced by the system

$$\begin{array}{l} y_{ijk} - x_{jk} + a_{ik} \geq 0, \quad k = 1, \dots, d, \\ y_{ijk} + x_{jk} - a_{ik} \geq 0, \quad k = 1, \dots, d, \\ y_{ijk}^{r} \leq \varsigma_{ijk}^{s} u_{ij}^{r-s}, \quad k = 1, \dots, d, \\ \omega_{i}^{\frac{r}{s}} \sum_{k=1}^{d} \varsigma_{ijk} \leq u_{ij}, \\ \varsigma_{ijk} \geq 0, \; \forall \; k = 1, \dots, d. \end{array}$$

Indeed, set $\rho = \frac{r}{r-s}$, then $\frac{1}{\rho} + \frac{s}{r} = 1$. Let $(\bar{x}_j, \bar{u}_{ij})$ fulfil the inequality $u_{ij} \ge \omega_i ||x_j - a_i||_{\tau}$. Then we have

$$\begin{split} \omega_{i} \|\bar{x}_{j} - a_{i}\|_{\tau} &\leq \bar{u}_{ij} \iff \omega_{i} \left(\sum_{k=1}^{d} |\bar{x}_{jk} - a_{ik}|^{\frac{r}{s}} \right)^{\bar{r}} \leq \bar{u}_{ij}^{\frac{s}{r}} \bar{u}_{ij}^{\frac{1}{p}} \\ &\iff \omega_{i} \left(\sum_{k=1}^{d} |\bar{x}_{jk} - a_{ik}|^{\frac{r}{s}} \bar{u}_{ij}^{\frac{r}{s}(-\frac{r-s}{r})} \right)^{\frac{s}{r}} \leq \bar{u}_{ij}^{\frac{s}{s}} \\ &\iff \omega_{i}^{\frac{r}{s}} \sum_{k=1}^{d} |\bar{x}_{jk} - a_{ik}|^{\frac{r}{s}} \bar{u}_{ij}^{-\frac{r-s}{s}} \leq \bar{u}_{ij} \end{split}$$
(24)

Then (24) holds if and only if $\exists \varsigma_{ij} \in \mathbb{R}^d$, $\varsigma_{ijk} \ge 0$, $\forall k = 1, ..., d$ such that

$$|\bar{x}_{jk} - a_{ik}|^{\frac{r}{s}} \bar{u}_{ij}^{-\frac{r-s}{s}} \leq \varsigma_{ijk}, \quad \text{satisfying} \quad \omega_i^{\frac{r}{s}} \sum_{k=1}^d \varsigma_{ijk} \leq \bar{u}_{ij},$$

or equivalently,

$$|\bar{x}_{jk} - a_{ik}|^r \le \varsigma_{ijk}^s \bar{u}_{ij}^{r-s}, \quad \omega_i^{\frac{r}{s}} \sum_{k=1}^d \varsigma_{ijk} \le \bar{u}_{ij}.$$
⁽²⁵⁾

Set $\bar{y}_{ijk} = |\bar{x}_{jk} - a_{ik}|$ and $\bar{\zeta}_{ijk} = |\bar{x}_{jk} - a_{ik}|^{\tau} \bar{u}_{ij}^{-1/\rho}$. Then, clearly $(\bar{x}_j, \bar{u}_{ij}, \bar{y}_{ij}, \bar{\zeta}_{ij})$ satisfies (6)–(9) and (14). Conversely, let $(\bar{x}_j, \bar{u}_{ij}, \bar{y}_{ij}, \bar{\zeta}_{ij})$ be a feasible solution of (6)–(9) and

(14). Then, $\bar{y}_{ijk} \ge |\bar{x}_{jk} - a_{ik}|$ for all *i*, *j* and by (8) $\bar{\varsigma}_{ijk} \ge \bar{y}_{ijk}^{(r_s)} u_{ij}^{-r_s} \ge |\bar{x}_{jk} - a_{jk}|^{\tau} \bar{u}_{ij}^{-r_s}$. Thus,

$$\omega_{i}^{\frac{r}{s}} \sum_{k=1}^{d} |\bar{x}_{jk} - a_{jk}|^{\frac{r}{s}} \bar{u}_{ij}^{-\frac{r-s}{s}} \le \omega_{i}^{\frac{r}{s}} \sum_{k=1}^{d} \bar{\zeta}_{ijk} \le \bar{u}_{ij},$$

which in turns implies that $\omega_i^{\frac{r}{3}} \sum_{k=1}^d |\bar{x}_{jk} - a_{jk}|^{\frac{r}{3}} \le \bar{u}_{ij}\bar{u}_{ij}^{\frac{r-s}{3}}$ and hence, $\omega_i \|\bar{x}_j - a_i\|_{\tau} \le \bar{u}_{ij}$. In the same way we prove that each inequality $t_{jj'} \ge \mu_{jj'} \|x_j - x_{j'}\|_{\tau}$, $j, j' = 1, \dots, p$ can be replaced by the system $z_{jj'k} - x_{jk} + x_{j'k} \ge 0$, $k = 1, \dots, d$, $z_{jj'k} + x_{jk} - x_{j'k} \ge 0$, $k = 1, \dots, d$,



Fig. 2. Demand points in Example 7 (filled circles) and solutions (triangles).

$$\begin{split} z_{jj'k}^{r} &\leq \xi_{jj'k}^{s} t_{jj'}^{r-s}, \ k = 1, \dots, d, \\ \mu_{jj'}^{\frac{r}{s}} &\sum_{k=1}^{d} \xi_{jj'k} \leq t_{jj'}, \\ &\xi_{ijk} \geq 0, \ \forall \ k = 1, \dots, d. \end{split}$$

Next, we observe that each one of the inequalities $y_{ijk}^r \leq \zeta_{ijk}^s u_{ij}^{r-s}$, $k = 1, \ldots, d$ (respectively $z_{jj'k}^r \leq \xi_{jj'k}^s t_{jj'}^{r-s}$, $k = 1, \ldots, d$) can be transformed, according to Lemma 2, into $4\log r$ linear matrix inequalities (respectively $4\log r$ linear matrix inequalities), being then exactly representable as second order cone constraints or semidefinite constraints.

Finally, it is straightforward to check Slater condition, for instance, for the system (23). Set $v_{ij} = 1$, $w_{lj} = \frac{3}{2}M\lambda_{lj}\max_i\omega_i$, $u_{ij} = \frac{3}{2}M\omega_i + 2$ with M > 0 large enough and $t_{jj'} = 2M\mu_{jj'} + 1$ for i, l = 1, ..., n and j, j' = 1, ..., p.

In the case that the considered norm is polyhedral (in particular if $\tau = 1$ or $\tau = +\infty$ in the theorem above), the above problem reduces to a standard linear problem and its number of variables and inequalities highly decreases.

As a consequence of Theorem 6 and Lemma 2, Problem (LOCOMF-NI) can be solved in polynomial time for any dimension *d*, by solving its reformulation as a SOCP problem (NIMFOMP_{λ}) (see Monteiro & Tsuchiya, 2000; Nesterov, Nemirovski, & Ye, 1994). The reader may note that this is an important step forward with respect to the already stated complexity results (see e.g. Rodríguez-Chía et al., 2000). There, it was proven that these problems are solvable in polynomial time in the plane (*d* = 2) and using polyhedral norms. Here we extend this complexity result to be true in any dimension *d* > 1 and for any polyhedral or ℓ_{τ} -norm with $\tau \in \mathbb{Q}$, $\tau \geq 1$.

Example 7. Consider a two-facility problem on the Euclidean plane with a set of 4 demand points given by $A = \{(9.46, 9.36), (8.93, 7.00), (2.20, 1.12), (1.33, 8.89)\}$ (*A* is subset of size 4 of the 50-cities data set from Eilon, Watson-Gandy, and Christofides (1971)), and (randomly generated-) lambda weights: $\lambda_{11} = 147.31$, $\lambda_{21} = 24.44$, $\lambda_{31} = 24.16$, $\lambda_{41} = 10.77$, $\lambda_{12} = 119.08)$, $\lambda_{22} = 0.56$, $\lambda_{32} = 0.00$, $\lambda_{42} = 0.00$, $\mu_{12} = 0.56$ and $w_i = 1$ for all i = 1, ..., n.

The problem to be solved can be written as

$$\min_{\substack{x_1, x_2 \in \mathbb{R}^2}} 147.31d_{(1)}(x_1) + 24.44d_{(2)}(x_1) + 24.16d_{(3)}(x_1) \\ + 10.77d_{(4)}(x_1) + 119.08d_{(1)}(x_2) + 0.56d_{(2)}(x_2) + 0.00d_{(3)}(x_2) \\ + 0.00d_{(4)}(x_2) + 0.56||x_1 - x_2||_2$$

We get as solution: $x_1^* = (5.24, 6.41)$ and $x_2^* = (5.61, 5.44)$, with objective value $f^* = 1704.55$. Fig. 2 shows the points and the

solutions of the problem. The reader may observe that in general optimal solutions of this problem do not collocate.

4. Single allocation multifacility location problems with ordered median objective functions

The structural difference of a single allocation multifacility location problem with those considered in the previous section rests on the fact that in the former each demand point shall be directed to a unique serving facility by means of a predetermined allocation rule (usually closest distance). This little difference makes the problem much more difficult since the convexity properties exhibited in the previous models are no longer valid and more sophisticated tools must be used to solve these problems.

In this framework, we are given a set $\{a_1, \ldots, a_n\} \subset \mathbb{R}^d$ endowed with a ℓ_{τ} -norm; and a feasible domain $\mathbf{K} = \{x \in \mathbb{R}^d : g_j(x) \ge 0, j = 1, \ldots, m\} \subset \mathbb{R}^d$, closed and semi-algebraic. The goal is to find p points $x_1, \ldots, x_p \in \mathbf{K} \subset \mathbb{R}^d$ minimizing some globalizing function of the shortest distances to the set of demand points.

The main feature and what distinguishes multifacility location problems from other general purpose optimization problems, is that the dependence of the decision variables is given throughout the norms to the demand points, i.e. $||x - a_i||_{\tau}$.

For the ease of presentation we have restricted ourselves to the particular case of pure location problems, namely $\tilde{f}_i(x) := \min_{j=1...p} ||x_j - a_i||_{\tau}$ which has attracted a lot of attention in the literature of Location Analysis. Needless to say that our methodology applies to more general forms of objective function, namely we could handle general rational functions of the distances as for instance by Blanco et al. (2013).

We shall define the dependence of the decision variables $x_1, \ldots, x_p \in \mathbb{R}^d$ via $t = (t_1, \ldots, t_n)$, where $t_i : \mathbb{R}^{pd} \mapsto \mathbb{R}$, $t_i(x_1, \ldots, x_p) := \min_j ||x_j - a_i||_{\tau}$, $i = 1, \ldots, n$. Therefore, the *i*th component of the ordered median objective function of our problems reads as

$$\begin{array}{cccc} \tilde{f}_i(x): & \mathbb{R}^{pd} & \mapsto & \mathbb{R} \\ & x = (x_1, \dots, x_p) & \mapsto & t_i := \min_{i=1} \{ \|x_j - a_i\|_\tau \}. \end{array}$$

Consider the following problem

$$\rho_{\lambda} := \min_{\mathbf{x}} \{ \sum_{i=1}^{n} \lambda_i \tilde{f}_{(i)}(\mathbf{x}) : \mathbf{x} = (x_1, \dots, x_p), \ \mathbf{x}_j \in \mathbf{K}, \ \forall j = 1, \dots, p \},$$
(LOCOMF)

where

- K ⊆ ℝ^d satisfies the Archimedean property. Without loss of generality we shall assume that we know M > 0 such that ||x_j||₂ ≤ M, for all j = 1, ..., p.
- $\tau := \frac{r}{s} \ge 1, r, s \in \mathbb{N}$ with gcd(r, s) = 1.
- $\lambda_{\ell} \geq 0$ for all $\ell = 1, \ldots, n$.

First of all, we observe that problem LOCOMF is well defined and that it has an optimal solution. Indeed, we are minimizing a continuous function over a compact set in \mathbb{R}^d . Thus, by Weierstrass theorem problem LOCOMF admits an optimal solution. On the other hand, unlike in the previous section uniqueness can not be easily obtained since, as proven by the following easy example, multiple solutions often occurs.

Example 8. Consider a two-facility problem on the Euclidean plane with set of demand points $A = \{ (0, 0), (0, 1), (1, 1), (1, 0) \}$. Assume that the λ -weights are $\lambda_{ij} = 1$ for all i = 1, ..., n, j = 1, ..., p. Then, any pair of points $x_1^* \in \{0\} \times [0, 1]$ and $x_2^* \in \{1\} \times [0, 1]$ is an optimal solution.

4.1. A mixed integer second order cone programming approach to solve LOCOMF

In this section, we present a tractable formulation of problem LOCOMF as a mixed integer nonlinear program with linear objective function. For each $i \in \{1, ..., n\}$, we set UB_i as a valid upper bound on the value of $\|\bar{x}_j - a_i\|_{\tau}$, $\bar{x}_j \in \mathbf{K}$.

We introduce the following auxiliary problem

$$\hat{\rho}_{\lambda} = \min \sum_{\ell=1}^{n} \lambda_{\ell} \theta_{\ell} \tag{MFOMP}_{\lambda}$$

s.t.
$$h_{il}^1 := t_i \le \theta_\ell + UB_i(1 - w_{i\ell}), \forall i = 1, ..., n, \ \ell = 1, ..., n,$$
 (26)

$$h_l^2 := \theta_\ell \ge \theta_{\ell+1}, \forall \ell = 1, \dots, n-1,$$
(27)

$$h_{ij}^3 := u_{ij} \le t_i + UB_i(1 - z_{ij}), \forall i = 1, \dots, n, \ j = 1, \dots, p,$$
(28)

$$h_{ijk}^4 \coloneqq v_{ijk} - x_{jk} + a_{ik} \ge 0, \forall i = 1, ..., n, \ j = 1, ..., p, \ k = 1, ..., d,$$
(29)

$$h_{ijk}^{5} \coloneqq v_{ijk} + x_{jk} - a_{ik} \ge 0, \forall i = 1, \dots, n, \ j = 1, \dots, p, \ k = 1, \dots, d,$$
(30)

$$h_{ijk}^{6} := v_{ijk}^{r} \le \zeta_{ijk}^{s} u_{ij}^{r-s}, \forall i = 1, \dots, n, \ j = 1, \dots, p, \ k = 1, \dots, d,$$
(31)

$$h_{ij}^7 := \sum_{k=1}^d \zeta_{ijk} \le u_{ij}, \forall i = 1, \dots, n, \ j = 1, \dots, p,$$
(32)

$$h_i^8 := \sum_{j=1}^p z_{ij} = 1, \, \forall i = 1, \dots, n,$$
 (33)

$$h_{\ell}^{9} := \sum_{i=1}^{n} w_{i\ell} = 1, \forall \ell = 1, \dots, n,$$
(34)

$$h_i^{10} := \sum_{l=1}^n w_{il} = 1, \forall i = 1, \dots, n,$$
 (35)

$$w_{i\ell} \in \{0, 1\}, \forall i, \ell = 1, ..., n,$$
 (36)

$$z_{ij} \in \{0, 1\}, \forall i = 1, \dots, n, j = 1, \dots, p,$$
(37)

$$\theta_{\ell}, t_i \in \mathbb{R}^+, \forall i, \ell = 1, \dots, n,$$
(38)

$$v_{ijk}, \ \zeta_{ijk}, \ u_{ij} \in \mathbb{R}^+, \ \forall i = 1, \dots, n, \ j = 1, \dots, p, \ k = 1, \dots, d,$$
 (39)

$$x_i \in \mathbf{K}, \forall j = 1, \dots, p. \tag{40}$$

With constraints (26) and (27), we enforce the variable θ_l to assume the value t_i that is sorted in the *l*th position of the vector *t*, while constraints (28)–(32) model the evaluation of $||x_j - a_i||_{\tau}$ for all *i* and *j*. Constraints (34)–(36) model permutations, and constraints (33) and (37) are introduced to model the allocation of element indexed by *i* to a unique index *j*. Therefore, putting all the above ingredients together we get that in the optimum $t_i = \min ||x_j - a_i||_{\tau}$.

Let $N := 3n + n^2 + n - 1 + np(3d + 2) = n^2 + 4n + np(3d + 2) - 1$ and denote by $\{h_1, \ldots, h_N\}$ the constraints (26)–(35) in the problem above. Let $\hat{\mathbf{K}}$ denote the feasible domain of Problem MFOMP₂.

Theorem 9. Let x be a feasible solution of LOCOMF then there exists a solution $(x, z, u, v, \zeta, w, t, \theta)$ for MFOMP_{λ} such that their objective values are equal. Conversely, if $(x, z, u, v, \zeta, w, t, \theta)$ is a feasible solution for MFOMP_{λ} then x is a feasible solution for LOCOMF. Furthermore, if K

satisfies Slater condition then the feasible region of the continuous relaxation of MFOMP_{λ} also satisfies Slater condition and $\rho_{\lambda} = \hat{\rho}_{\lambda}$.

Proof. Let $\bar{x} = (\bar{x}_1, ..., \bar{x}_p)$ be a feasible solution of LOCOMF. Then, it satisfies $\bar{x}_j \in \mathbf{K}$, for all j = 1, ..., p. Let $u_{ij} = \|\bar{x}_j - a_i\|_{\tau}$, based in (24) and (25), $\|\bar{x}_j - a_i\|_{\tau}$ can be represented by

$$\begin{cases} \nu_{ijk} = |\bar{x}_{jk} - a_{ik}|, \\ \nu_{ijk}^{r} = \zeta_{ijk}^{s} u_{ij}^{r-s}, \\ \sum_{k=1}^{d} \zeta_{ijk} = u_{ij}, \\ \zeta \ge 0. \end{cases}$$

For i = 1, ..., n, j = 1, ..., p and k = 1, ..., d, we denote by

$$t_{i} = \min_{\ell} \|\bar{x}_{\ell} - a_{i}\|_{\tau} \quad \text{and} \quad z_{ij} = \begin{cases} 1, & \text{if } \min_{l} \|\bar{x}_{l} - a_{i}\|_{\tau} \\ &= \|\bar{x}_{j} - a_{i}\|_{\tau}, \\ 0, & \text{otherwise.} \end{cases}$$

Observe that if it would exist $j' \in \{1, ..., p\}$ such that $j' \neq j$ and $\min_{l} \|\bar{x}_{l} - a_{i}\|_{\tau} = \|\bar{x}_{j'} - a_{i}\|_{\tau}$ then we can choose arbitrarily any of them, because a client can be assigned to only one facility.

These values clearly satisfy constraints (28)–(32) and (37)–(39).

Besides, let σ be the permutation of (1, ..., n) such that $t_{\sigma(1)} \ge ... \ge t_{\sigma(n)}$. Take,

$$w_{il} = \begin{cases} 1, & \text{if } i = \sigma(l), \\ 0, & \text{otherwize;} \end{cases} \text{ and } \theta_l = t_{\sigma(l)} \end{cases}$$

Then the constraints (26) and (27) are also satisfied. Clearly, $\sum_{\ell=1}^{n} \lambda_{\ell} \theta_{\ell} = \sum_{i=1}^{n} \lambda_{i} \tilde{f}_{(i)}(x)$.

Conversely, if $(\bar{x}, \bar{z}, \bar{u}, \bar{v}, \bar{\zeta}, \bar{w}, \bar{t}, \bar{\theta})$ is a feasible solution of MFOMP_{λ} then, clearly $\bar{x}_i \in \mathbf{K}$ and \bar{x} is a feasible point of LOCOMF.

Consider the continuous relaxation of problem (MFOMP_{λ}). Suppose that **K** satisfies Slater condition. Take x_j for all j = 1, ..., p, in the interior of **K**. Set $\theta = 4M + \frac{1}{\ell}$, $t_i = 3M$ and $u_{ij} = 2M$ for ℓ , i = 1, ..., n and j = 1, ..., p with M > 0 large enough. Then, for any z_{ij} , $w_{i\ell} \in [0, 1]$ we get that the set of inequality constraints satisfies

$$\begin{cases} \theta_{\ell} - t_i + UB_i(1 - w_{i\ell}) > 0, & i = 1, \dots, n, \ \ell = 1, \dots, n, \\ \theta_{\ell} > \theta_{\ell+1}, & \ell = 1, \dots, n, \\ t_i - u_{ij} + UB_i(1 - z_{ij}) > 0, & \forall i = 1, \dots, n, \ j = 1, \dots, p, \\ u_{ij} > \|x_j - a_i\|_{\tau}, & \forall i = 1, \dots, n, \ j = 1, \dots, p. \end{cases}$$

This proves that the continuous relaxation of MFOMP_λ satisfies Slater condition.

By the above arguments, optimal solutions and optimal values of both formulations coincide.

Example 10. To illustrate the applicability of the above formulation, in the following we have applied it to an instance of 10 demand points extracted from the 50-points data set by Eilon et al. (1971):

(9.46, 9.36), (7.43, 1.61), (6.27, 3.66), (5.00, 9.00), (2.83, 9.88), (2.20, 1.12), (1.90, 8.35), (1.68, 6.45), (1.24, 6.69), (0.75, 4.98).

For p = 3, $\tau = \frac{7}{5}$ and $\lambda = (2.25, 1.70, 1.14, 1.11, 1.06, 1.03, 1.01, 1.01, 1.00, 1.00)$, we get the solutions $x_1^* = (6.19, 1.58)$, $x_2^* = (5.00, 9.36)$, and $x_3^* = (1.44, 6.55)$, with optimal objective value $f^* = 30.1460$. Fig. 3 shows the demand points (filled dots), solutions (filled triangles) and the allocation of the demand points to the facilities (dashed lines).

An interesting observation that follows from Problem (MFOMP_{λ}) is that the unconstrained version of the location problem is an appealing mixed integer nonlinear program. Observe that the only set of nonlinear constraints that appear in (MFOMP_{λ}) is (31). However, (31) can be written equivalently as a polynomial number of second order cone constraints, according to Lemma 2. This way, Problem

 $(MFOMP_{\lambda})$ becomes a mixed integer nonlinear program with lineal objective function and only linear and second order cone constraints, although with two sets of binary variables, namely *w* and *z*. Nevertheless, there are nowadays general purpose solvers, as Gurobi, Cplex or Xpress, that implements exact B&B algorithms for this type of problems and that are rather efficient.

Remark 11. Assume that the hypothesis of Theorem 9 are satisfied. In addition, any of the following conditions hold

- 1. $g_i(x)$ are concave for i = 1, ..., m and $-\sum_{i=1}^m \mu_i \nabla^2 g_i(x) > 0$ for each dual pair (x, μ) of the problem of minimizing any linear functional $c^t x$ on **K** (*Positive Definite Lagrange Hessian* (PDLH)).
- 2. $g_i(x)$ are sos-concave on **K** for i = 1, ..., m or $g_i(x)$ are concave on **K** and strictly concave on the boundary of **K** where they vanish, i.e. $\partial \mathbf{K} \cap \partial \{x \in \mathbb{R}^d : g_i(x) = 0\}$, for all i = 1, ..., m.
- 3. $g_i(x)$ are strictly quasi-concave on **K** for i = 1, ..., m.

Then, applying (Blanco et al., 2014b, Theorem 4) to the formulation of Problem (MFOMP_{λ}), we get that there exists a constructive finite dimension embedding, which only depends on τ and g_i , i = 1, ..., m, such that the problem is mixed-integer SDP representable.

From the above observation to solve Problem MFOMP_{λ} efficiently we have combined a branch-and-bound approach over a mixed integer nonlinear program. In that approach, one may provide two types of lower bounds in the nodes of the branching tree: a continuous relaxation and a SDP relaxation based on a hierarchy of SDP "*a la Lasserre*". Clearly, the first type of bounds are only possible if in each node of the tree the continuous relaxation of MFOMP_{λ} satisfies Slater condition. This is ensured in Theorem 9.

The consequence of the above transformation is that one can easily put this family of problems in commercial solvers and then to get solutions without going to painful *ad hoc* implementations that may be problem dependent. We illustrate this approach in our computational experiments in Section 5.

Finally, we conclude this section providing some results concerning the second type of lower bounds, based on a SDP hierarchy, that can be used to approximate to any degree of accuracy the solution of the problem as well as within the branch and bound framework at each node of the branching tree.

Let $\mathbf{y} = (y_{\alpha})$ be a real sequence indexed in the monomial basis $(x^{\beta}z^{\eta}u^{\gamma}v^{\delta}\zeta^{\kappa}w^{\psi}t^{\tau}\theta^{\vartheta})$ of $\mathbb{R}[x, z, u, v, \zeta, w, t, \theta]$ (with $\alpha = (\beta, \eta, \gamma, \delta, \kappa, \tau, \psi, \vartheta) \in \mathbb{N}^{pd} \times \mathbb{N}^{np} \times \mathbb{N}^{npd} \times \mathbb{N}^{n^2} \times \mathbb{N}^n \times \mathbb{N}^n \times \mathbb{N}^{nd}$). Denote by $nv_1 = d + np(d+2) + n^2 + 2n + npd$ the number of variables in the extended formulation of the problem.

Let
$$h_0(\theta) := \sum_{\ell=1}^m \lambda_\ell \theta_\ell$$
, and denote $\xi_j := \lceil (\deg g_j)/2 \rceil$ and $\nu_j :=$

 $\lceil (\deg h_j)/2 \rceil$, where $\{g_1, \ldots, g_{n_{\mathbf{K}}}\}$, and $\{h_1, \ldots, h_N\}$ are, respectively, the polynomial constraints that define **K** and $\hat{\mathbf{K}} \setminus \mathbf{K}$ in MFOMP_{λ}. For $r \ge r_0 := \max\{\max_{k=1,\ldots,n_{\mathbf{K}}} \xi_k, \max_{j=0,\ldots,N} \nu_j\}$, introduce the hierarchy of semidefinite programs:

$$\inf_{\mathbf{y}} \quad L_{\mathbf{y}}(p_{\lambda})$$

s.t.
$$M_r(\mathbf{y}) \succeq 0$$
,
 $M_{r-\xi_k}(g_k, \mathbf{y}) \succeq 0$, $k = 1, \dots, n_{\mathbf{K}}$, $Q1_r$
 $M_{r-\nu_j}(h_j, \mathbf{y})(\stackrel{\succ}{=})0$, $j = 1, \dots, N$,
 $y_0 = 1$,

with optimal value denoted $\inf \mathbf{Q1}_r$ (and $\min \mathbf{Q1}_r$ if the infimum is attained). The reader may observe that some of the localizing constraints $M_{r-\nu_j}(h_j, \mathbf{y})$ in $\mathbf{Q1}_r$ are linear matrix equations since the families of constraints (33)–(35) contain several equations. This is indicated in $\mathbf{Q1}_r$ by our notation $M_{r-\nu_j}(h_j, \mathbf{y})(\succeq)$. Next, based on Proposition 1 and (Lasserre, 2009b, Theorem 6.1) we can state the following result.



Fig. 3. Demand points in Example 10 (filled circles), solutions (triangles) and allocation of demand points to facilities (dashed lines).

Theorem 12. Let $\hat{\mathbf{K}} \subset \mathbb{R}^{n\nu_1}$ (compact) be the feasible domain of Problem MFOMP_{λ}. Let inf $\mathbf{Q1}_r$ be the optimal value of the semidefinite program $\mathbf{Q1}_r$. Then, with the notation above:

- (a) $\inf \mathbf{Q1}_r \uparrow \rho_\lambda \text{ as } r \to \infty$.
- (b) Let \mathbf{y}^r be an optimal solution of the SDP relaxation $Q1_r$. If

rank
$$M_r(\mathbf{y}^r) = \operatorname{rank} M_{r-r_0}(\mathbf{y}^r) = \varphi$$

then $\min \mathbf{Q1}_r = \rho_{\lambda}$ and one may extract φ points $(x_1^*(k), \ldots, x_p^*(k), z^*(k), u^*(k), v^*(k), \zeta^*(k), w^*(k), t^*(k), \theta^*(k))_{k=1}^{\varphi} \subset \mathbf{\hat{K}}$, all global minimizers of the MFOMP_{λ} problem.

One of the drawbacks of using hierarchies of SDP problems to approximate the solution of our original multifacility location problem $(MFOMP_{\lambda})$ is the dimension of the SDP objects that must be used when the relaxation order increases. The size of the matrices to be considered in the SDP problems that have to be solved grows exponentially with the relaxation order. Significant reductions in the sizes of the problems to be solved can be performed by exploiting the sparsity and symmetry that appear in the location problem. On the one hand, at times the number of variables that appear in the polynomial constraints of $(MFOMP_{\lambda})$ can be separated in blocks so that only some of them appear together in some constraints. In those cases the SDP variables to be used can be simplified and thus, the sizes of the moment and localizing matrices are dramatically reduced. The application of this result requires the so call running intersection property (see Lasserre, 2006) that easily applies in this case. On the other hand, the invariance of this problem under the action of the symmetric group applied to the indices of the facilities (*j*) allows to adapt the results by Riener, Theobald, Andrén, and Lasserre (2013) to reduce the dimensionality of the matrices that appear in the hierarchy of SDP $(Q1_r)$ that converge to the optimal solution. The interested reader can find all the details of the application of the dimensionality reduction based on sparsity and symmetry for multifacility location problems by Blanco, Puerto, and El-Haj Ben-Ali (2014a). Combining both approaches helps to solve some problems.

5. Computational experiments

We have performed a series of computational experiments to show the efficiency of all the proposed formulations and approaches. The SOCP (LOCOMF-NI) and the mixed integer SOCP (MFOMP_{λ}) formulations have been coded in Gurobi 5.6 and executed in a PC with an Intel Core i7 processor at 2x 2.40 GHz and 4 GB of RAM. We have applied our formulations to the well-known 50-points data set by Eilon et al. (1971) by considering different number of facilities, norms

Table 1								
CPU	running	times	for	non-interchangeable	multifacility			
problem for the Eilon–Watson–Christofides 50-point data set.								

р	τ	τ							
	1.5	2	3						
2	2.5095	2.1157	3.7470						
5	12.7794	6.5161	9.8130						
10	29.1873	10.5726	19.5455						
15	49.4854	19.1129	40.4506						
30	148.7449	40.5635	85.5676						

and weights. In particular we have considered the number of facilities to be located, *p*, ranging in {2, 5, 10, 15, 30} and τ (the norm) in { $\frac{3}{2}$, 2, 3}. We have run both the unconstrained non-interchangeable LOCOMF-NI model and the single-allocation model MFOMP_{λ}. For the non-interchangeable model we have considered random λ and μ weights (fulfiling the conditions described in (3)). Table 1 reports our results on these experiments. There, we report the average CPU times of 5 different random instances for each problem (when fixing *p* and τ).

For general single-allocation multifacility problems, we have consider three of the most classical multifacility models that fit within the ordered median formulation when particular λ weights are chosen: *p*-median problem, *p*-center problem and *p*-*k*-centrum problem (with k = 0.5 n = 25). In Table 2 we report the overall CPU times needed to solve the problems and the optimal solutions provided by Gurobi (f^*). These optimal values are important to test accuracy of solutions for future authors that wish to compare new approaches to solve these problems.

We observe that the bottleneck for solving (MFOMP_{λ}) is the existence of binary variables (*z* and *w*), since otherwise it would be solvable in polynomial time. Due to the high running time, optimality could not be ensured for some instances. In order to ease the resolution some improvements can be done via some preprocessing and fixing binary variables based on a geometric branch and bound phase. This preprocessing is effective whenever the dimension of the space of variables in low, namely *d* = 2, 3.

Finally, to illustrate the use of the hierarchy of problems $Q1_r$, we have applied to the same data set by Eilon et al. (1971) the SDP relaxation $Q1_2$ for *p*-median problems with Euclidean distances. Implementation was done in SDPT3. The results are detailed in Table 3. The reader can observe that the results obtained with this approach (in terms of accuracy) are good although we could no certify optimality since the rank condition was not satisfied. Moreover, the running times are considerably higher (by several orders of magnitude) than those needed for similar problems solved with the MISOCO implementations in Gurobi.

Table 2

Computational results for *p*-median, *p*-center and *p*-25-centrum problems for the 50-points data set by Eilon et al. (1971).

р	τ	p-median		<i>p</i> -center		p-25-centrum	
		CPUTime	<i>f</i> *	CPUTime	<i>f</i> *	CPUTime	f^*
2	1.5	22.31	150.955	1.03	4.9452	10.08	100.8474
	2	1.13	135.5222	0.28	4.8209	0.38	95.0892
	3	23.68	130.8560	13.51	4.7880	139.03	89.0238
5	1.5	55.28	78.6074	3.73	2.8831	33.09	53.4995
	2	12.49	72.2369	5.37	2.6610	7.61	49.6932
	3	125.10	68.1791	2.87	2.5094	18.23	46.9844
10	1.5	5.36	45.0525	2.66	1.6929	68.36	30.7137
	2	2.31	41.6851	5.3	1.6113	17.93	28.9017
	3	4.76	39.7222	55.76	1.5950	225.64	27.5376
15	1.5	6.70	30.0543	9.44	1.1139	49.92	22.4165
	2	43.91	27.6282	0.62	1.0717	11.26	20.6536
	3	150.99	26.6047	50.08	1.0530	244.59	20.8544
30	1.5	14.45	9.9488	74.43	1.0080	202.54	9.0806
	2	4.81	8.7963	1.53	0.9192	5.29	8.5216
	3	198.78	8.6995	57.37	0.8508	287.90	8.0016

Table 3

Second relaxations in the Lasserrs's hierarchy for Euclidean *p*-median problems for the data set by Eilon et al. (1971).

ар

6. Conclusions

We propose a novel, effective mixed integer nonlinear programming formulation for the continuous multifacility ordered median location problem for any ℓ_{τ} -norm and in any dimension. This formulation provides a unified approach for dealing with a broad family of multifacility location problems which up to now were usually solved only for some special cases and usually in low dimension. We also provide lower bounds for the nodes of the branching tree based on a hierarchy of SDP relaxations "*a la Lasserre*". A simpler class of multifacility (multiple-allocation) location problems is also analyzed for which optimal solutions are obtained by a continuous SOCP formulation.

References

- Ahuja, R. K., Magnanti, T. L., & Orlin, J. B. (1993). Network flows. Theory, algorithms, and applications. Prentice Hall.
- Albareda, M., Hinojosa, Y., & Puerto, J. (2015). The reliable p-median problem with atfacility service. European Journal of Operational Research, 245(3), 656–666.
- Alizadeh, F., & Goldfarb, D. (2003). Second order cone programming. Mathematical Programming B, 95, 3–51.
- Anjos, M. F., & Lasserre, J. B. (2012). Handbook on semidefinite, conic and polynomial optimization. Springer.
- Blanco, V., El-Haj Ben-Ali, S., & Puerto, J. (2013). Minimizing ordered weighted averaging of rational functions with applications to continuous location. *Computers & Operations Research*, 40, 1448–1460.
- Blanco, V., Puerto, J. El-Haj Ben-Ali, S. (2014a). Algorithms and dimensionality reductions for continuous multifacility ordered median location problems. http: //arxiv.org/abs/1401.0817.
- Blanco, V., Puerto, J., & El-Haj Ben-Ali, S. (2014b). Revisiting several problems and algorithms in continuous location with ℓ_p norms. *Computational Optimization and Applications*, 58(3), 563–595.
- Boland, N., Dominguez-Marin, P., Nickel, S., & Puerto, J. (2006). Exact procedures for solving the discrete ordered median problem. *Computers & Operations Research*, 33, 3270–3300.
- Boyd, S., & Vanderberghe, L. (2004). Convex optimization. Cambridge University Press.

Brimberg, J., Drezner, Z., Mladenović, N., & Salhi, S. (2014). A new local search for continuous location problems. *European Journal of Operational Research*, 232(2), 256–265.Brimberg, J., Hansen, P., Mladenovic, N., & Taillard, E. D. (2000). Improvement and com-

- Brimberg, J., Hansen, P., Mladenovic, N., & Taillard, E. D. (2000). Improvement and comparison of heuristics for solving the uncapacitated multisource weber problem. *Operations Research*, 48, 444–460.
- Carrizosa, E., & Puerto, J. (1995). A discretizing algorithm for location problems. *European Journal of Operational Research*, 80(1), 166–174.
- Drezner, Z., & Hamacher, H. W. (2002). Facility location: applications and theory. Springer.
- Eilon, S., Watson-Gandy, C., & Christofides, N. (1971). Distribution management: mathematical modeling and practical analysis. London: Griffin.
- Kalcsics, J., Nickel, S., Pozo, M. A., Puerto, J., & Rodriguez-Chia, A. M. (2014). The multicriteria p-facility median location problem on networks. *European Journal of Operational Research*, 235(3), 484–493.
- Kalcsics, J., Nickel, S., Puerto, J., & Rodríguez-Chía, A. M. (2010). The ordered capacitated facility location problem. *Top*, 18(1), 203–222.
- Kariv, O., & Hakimi, S. L. (1979). An algorithmic approach to network location problems i. SIAM Journal of Applied Mathematics, 37, 513–538.
- Lasserre, J. B. (2001). Global optimization with polynomials and the problem of moments. SIAM Journal on Optimization, 11, 796–817.
- Lasserre, J. B. (2006). Convergent sdp-relaxations in polynomial optimization with sparsity. SIAM Journal on Optimization, 17, 822–843.
- Lasserre, J. B. (2009a). Moments and sums of squares for polynomial optimization and related problems. *Journal of Global Optimization*, 45, 39–61.
- Lasserre, J. B. (2009b). Moments, positive polynomials and their applications. Imperial College Press.
- Lobo, M. S., Vandenberghe, L., Boyd, S., & Lebret, H. (1998). Applications of second-order cone programming. *Linear Algebra and its Applications*, 284(1–3), 193–228.
- Marín, A., Nickel, S., Puerto, J., & Velten, S. (2009). A flexible model and efficient solution strategies for discrete location problems. *Discrete Applied Mathematics*, 157(5), 128– 1145.
- Michelot, C. (1987). Localization in multifacility location theory. European Journal of Operational Research, 31(2), 177–184.
- Monteiro, R. D. C., & Tsuchiya, T. (2000). Polynomial convergence of primal-dual algorithms for the second-order cone program based on the mz-family of directions. *Mathematical Programming*, 88, 61–83.

Nesterov, Y., Nemirovski, A., & Ye, Y. (1994). Interior-point polynomial algorithms in convex programming. *SIAM: Studies in Applied and Numerical Mathematics*, 13.

Nickel, S., & Puerto, J. (2005). Facility location - a unified approach. Springer. Puerto, J., D. Pérez-Brito, D., & Garcia-González, C. G. (2014). A modified variable neigh-

- borhood search for the discrete ordered median problem. European Journal of Operational Research, 234(1), 61–76.
- Puerto, J., & Rodriguez-Chia, A. M. (2011). On the structure of the solution set for the single facility location problem with average distances. *Mathematical Programming*, 128, 373–401.
- Putinar, M. (1993). Positive polynomials on compact semi-algebraic sets. Indiana University Mathematics Journal, 42, 969–984.
- Riener, C., Theobald, T., Andrén, L. J., & Lasserre, J. B. (2013). Exploiting Symmetries in SDP-Relaxations for polynomial optimization. *Mathematics of Operations Research*, 3, 122–141.
- Rodríguez-Chía, A. M., Nickel, S., Puerto, J., & Fernández, F. R. (2000). A flexible approach to location problems. *Mathematical Methods of Operations Research*, 51(1), 69–89.
- Wolkowicz, H., Saigal, R., & Vandenberghe, L. (2000). Handbook of semidefinite programming. Kluwer Academic Publishers.