Contents lists available at ScienceDirect



European Journal of Operational Research

journal homepage: www.elsevier.com/locate/ejor

# Ordered Weighted Average optimization in Multiobjective Spanning Tree Problem



CrossMark

UROPEAN JOURNAL PERATIONAL RESEA

## Elena Fernández<sup>a</sup>, Miguel A. Pozo<sup>b</sup>, Justo Puerto<sup>b,\*</sup>, Andrea Scozzari<sup>c</sup>

<sup>a</sup> Department of Statistics and Operational Research, Universitat Politècnica de Catalunya, Barcelona Tech. Jordi Girona 1-3, Barcelona 08034, Spain <sup>b</sup> IMUS and Faculty of Mathematics, Universidad de Sevilla, Seville 41012, Spain

<sup>c</sup> Facoltà di Economia, Università degli Studi Niccolò Cusano, Roma - 00166 Roma, Italy

#### ARTICLE INFO

Article history: Received 2 July 2015 Accepted 12 October 2016 Available online 18 October 2016

Keywords: Combinatorial optimization Multiobjective optimization Ordered median Ordered Weighted Average Spanning trees

#### ABSTRACT

Multiobjective Spanning Tree Problems are studied in this paper. The ordered median objective function is used as an averaging operator to aggregate the vector of objective values of feasible solutions. This leads to the Ordered Weighted Average Spanning Tree Problem, a nonlinear combinatorial optimization problem. Different mixed integer linear programs are proposed, based on the most relevant minimum cost spanning tree models in the literature. These formulations are analyzed and several enhancements presented. Their empirical performance is tested over a set of randomly generated benchmark instances. The results of the computational experiments show that the choice of an appropriate formulation allows to solve larger instances with more objectives than those previously solved in the literature.

© 2016 Elsevier B.V. All rights reserved.

## 1. Introduction

Optimization problems related to spanning trees, or simply Spanning Tree Problems are among the core problems in combinatorial optimization. On the one hand, the combinatorial object that represents spanning trees has important structural properties. On the other hand, from a practitioner point of view, spanning trees are found in a wide range of applications in many fields (e.g. computer networks design, telecommunications networks, transportation, etc.). Furthermore, they often appear as subproblems of other more complex optimization problems.

The most relevant property of trees is their matroid structure. This implies that the basic problem of finding a minimum cost spanning tree, can be solved efficiently (Kruskal, 1956; Prim, 1957). This also implies that formulations with the integrality property can be obtained, which allow to solve the minimum cost Spanning Tree Problem (STP) with linear programing tools. However, these good features can be lost for several reasons. For instance, when the objective function does not preserve Gale optimality, i.e., it is not monotonic on the edges costs (Fernández, Pozo, & Puerto, 2014; Lawler, 1966), as it happens in the Optimum Communication Spanning Tree Problem (Hu, 1974). The reader may refer to Landete and Marín (2014) for a description of alternative objective func-

tions for STP. An optimization STP also becomes a hard problem when several objectives are considered simultaneously (Ehrgott, 2005). In such cases, no efficient combinatorial algorithm is known so the choice of an appropriate mathematical programing representation of the combinatorial object may become crucial. In this sense, formulations for STP with good properties can be outperformed by other formulations in the new environment.

From a different point of view, in the multiobjective case, it is widely accepted that the use of order and aggregation functions may yield compromise solutions for the different criteria. The literature includes many works on this area related to combinatorial optimization. Some examples, among many others, include minimax problems (Hansen, 1979; Schrijver, 1983), combining minisum and minimax (Averbakh & Berman, 1995; Hansen & Labbé, 1988; Hansen, Labbé, & Thisse, 1991; Minoux, 1989; Punnen, Nair, & Aneja, 1995; Tamir, Puerto, & Perez-Brito, 2002), kcentrum optimization (Garfinkel, Fernández, & Lowe, 2006; Kalcsics, Nickel, Puerto, & Tamir, 2002; Punnen, 1992; Slater, 1978a; 1978b; Tamir, 2000), lexicographic optimization (Calvete & Mateo, 1998; Croce, Paschos, & Tsoukias, 1999), kth best solutions (Lawler, 1972; Martello, Pulleyblank, Toth, & de Werra, 1984; Pascoal, Captivo, & Clímaco, 2003; Yen, 1971), most uniform solutions (Galil & Schieber, 1998; López de los Mozos, Mesa, & Puerto, 2008), minimum-envy solutions (Espejo, Marin, Puerto, & Rodríguez-Chía, 2009), solutions with minimum deviation (Gupta, Gupta, & Bector, 1990), regret solutions (Averbakh, 2001; Conde, 2004; Puerto & Rodríguez-Chía, 2003), equity measures (Gupta & Punnen, 1988; Mesa, Puerto, & Tamir, 2003; López de los Mozos et al., 2008;

<sup>\*</sup> Corresponding author. Fax: +34 954622800.

*E-mail addresses:* e.fernandez@upc.edu (E. Fernández), miguelpozo@us.es (M.A. Pozo), puerto@us.es, justo.puerto@gmail.com (J. Puerto), andrea.scozzari@ unicusano.it (A. Scozzari).

Punnen & Aneja, 1997), discrete ordered median location problems (Boland, Domínguez-Marín, Nickel, & Puerto, 2006; Marín, Nickel, Puerto, & Velten, 2009; Puerto, 2008; Puerto & Rodríguez-Chía, 2015; Puerto & Tamir, 2005), Ordered Weighted Average objectives (Fernández, Pozo, & Puerto, 2013; 2014; Galand & Spanjaard, 2012), and covering objectives (Balas & Padberg, 1972; Breuer, 1970; Christofides & Korman, 1974; Kelly, 1944; Lawler, 1966). This paper elaborates on a specific form of aggregation criterion in multicriteria optimization, called Ordered Weighted Average operator (OWA). It is well known that this family parameterizes the aggregation function used when the decision-maker seeks the simultaneous satisfaction of all the criteria, to the case when the individual satisfaction of any of the criteria is sought. This fact is particularly relevant, due to its generality, as it includes as particular cases most of the above mentioned operators. This observation connects our results with the multicriteria literature and has been made already explicit in Fernández et al. (2014).

Multiobjective STPs have already been studied by some authors, mostly for the biobjective case (see Hamacher & Ruhe, 1994; Andersen, Jörnsten, & Lind, 1996; Ramos, Alonso, Sicilia, & González, 1998; Sourd & Spanjaard, 2008; Steiner & Radzik, 2008). In this paper we address the Multiobjective STP under the perspective of the OWA operator for a general number of objectives. This problem will be referred to as OWA Spanning Tree Problem (OWASTP). In OWASTP the optimality of traditional combinatorial algorithms is no longer guaranteed. Furthermore, formulations adapted from good STP formulations lose the integrality property. Thus alternative formulations that originally do not exhibit such good properties, may now outperform them. In Galand, Perny, and Spanjaard (2010) OWASTP was addressed using Choquet optimization and Galand and Spanjaard (2012) presented a first ordered median Mixed Integer Linear Programing (MILP) formulation. Our goal in this paper is to exploit properties of alternative formulations for OWASTP. As we will see, an appropriate formulation allows us to solve larger instances and with more objectives than those previously solved in the literature (Galand & Spanjaard, 2012), with up to 100 nodes and 10 objective functions. The contributions of this paper are (1) to provide new formulations for OWASTP combining appropriate formulations for STP and for OWA problem; (2) to prove a new complexity result showing that OWASTP is NPcomplete even for cactus graphs and two objectives; (3) to establish a theoretical and empirical comparison between the new formulations and previous existing ones; and, (4) to provide reinforcements that together with the new OWASTP formulations are able to outperform previous results in the literature.

The structure of the paper is the following. In Section 2 we formally define OWASTP and prove our new complexity result. Section 3 presents the catalogue of STP formulations that we study for OWASTP. One such formulation has already been used in Galand and Spanjaard (2012). We will use it as a reference for the alternative formulations that we present. The empirical performance of the resulting OWASTP formulations is analyzed in Section 4, where we present extensive numerical results and a comparison with existing ones. Finally, some conclusions are summarized in Section 5.

## 2. Problem definition

The Ordered Weighted Average operator is defined over a feasible set  $Q \subseteq \mathbb{R}^m$ . Let  $C \in \mathbb{R}^{p \times m}$  be a given matrix, whose rows, denoted by  $C^i$ , are associated with the cost vectors of p objective functions. The index set for the rows of C is denoted by  $P = \{1, \ldots, p\}$ . Let also  $\omega \in \mathbb{R}^{p+}$  denote a vector of non-negative weights. For  $x \in Q$ , the vector  $y = Cx \in \mathbb{R}^p$  is referred to as the outcome vector relative to C. In the following we assume y = Cx, with  $x \in Q$ . For a given y, let  $\sigma$  be a permutation of the indices of  $i \in$  *P* such that  $y_{\sigma_1} \ge \cdots \ge y_{\sigma_p}$ . Feasible solutions  $x \in Q$  are evaluated with an operator defined as  $OWA_{(C,\omega)}(x) = \omega^t y_{\sigma}$ . The OWA optimization Problem (OWAP) is to find  $x \in Q$  of minimum value with respect to the above operator, that is

## OWAP: $\min_{x \in O} OWA_{(C,\omega)}(x)$

The OWA is a very general operator, which has as particular cases well-known objective functions namely the Ordered Median Objective and the Vector Assignment Ordered Median (see Fernández et al., 2014). In addition, the OWA operator allows to model various aggregation functions according to the vector of weights *w* (see e.g., Ogryczak & Olender, 2012). Some examples are the minimum, maximum, median, center or *k*-centrum functions. Therefore, the selection of non/monotonic or non/symmetric *w*-weights is directly connected with the problem structure and thus with its complexity (Kasperski & Zielinski, 2015).

As defined, the OWA operator is indeed not linear. Moreover, in general, it is not convex either. For the case of monotonic weights, its convexity is known (Kalcsics, Nickel, & Puerto, 2003; Puerto & Rodríguez-Chía, 2015) and some elegant linearization of OWA functions have been proposed in the literature (see e.g., Ogryczak & Sliwinski, 2003; Ogryczak & Tamir, 2003). Depending on the type of monotonicity, the problems are simpler (with decreasing weights in the case of minimization) or harder (with increasing weights in the case of minimization). In this paper we focus on OWASTP with arbitrary weights. Two well-known particular cases of the OWA operator with arbitrary weights are the Hurwicz criterion (Hurwicz, 1951) defined as  $\alpha \max_{i \in P} y_i + (1 - \alpha) \min_{i \in P} y_i$  and the *k*-trimmed mean defined as  $\sum_{i=k+1}^{p-k} (p-2k)^{-1}y_i$ . These criteria are of special interest for being non-monotonic and non-convex (Grzybowski, Nickel, Pallaschke, & Urbanski, 2011, Puerto & Tamir, 2005) and have already been considered when analyzing the behavior of OWA operators in multiobjective optimization (see e.g., Galand & Spanjaard, 2012).

OWASTP is defined as follows. Let G = (V, E) be an undirected connected graph with set of nodes V, |V| = n, and set of edges E, |E| = m. In the following we assume that G contains at least one cycle, that is m > n - 1, as otherwise the problem becomes trivial. A spanning tree of G is a subgraph T = (V, E') where  $E' \subset E$  is a minimal set of edges connecting the set of nodes V. Let T denote the set of spanning trees defined on G. Then, OWASTP can be defined as

OWASTP: 
$$\min_{x \in T} OWA_{(C,\omega)}(x)$$
.

**Example 1.** Consider the graph G = (N, E) depicted in Fig. 1(a) and the 3-cost vectors on *E*, whose values are represented next to each edge. The optimal solution to OWASTP with  $\omega^t = (0.4, 0, 0.6)$  is depicted in Fig. 1(b) and has a value of 8.8. When the weights are  $\omega^t = (0.8, 0, 0.2)$  the optimal OWASTP value is 10.4, corresponding to the tree depicted in Fig. 1(c).

OWASTP is known to be NP-hard on general graphs since, for a weight vector  $w_1 = 1$  and  $w_i = 0$ , i = 2, ..., p, the problem reduces to the bicriteria Min–Max Spanning tree which is NP-hard on general graphs (Hamacher & Ruhe, 1994). This complexity result can be strengthened following (Yu, 1998), where the author proved that the bicriteria Min–Max Spanning Tree Problem is NPhard on grid graphs and p = 2. Consequently, OWASTP is NP-hard in this special class of graphs, as well. However, we can further refine the latter complexity by showing that OWASTP remains NPhard on the very special class of *cactus* graphs, for p = 2, and assuming a strictly decreasing weights vector.

Given a connected graph G = (V, E), a node  $v \in V$  is a cut node if after removing v and all edges incident to it the remaining graph is no longer connected. A graph without cut vertices is called



**Fig. 1.** Graph with edge costs (a) and OWASTP solutions for  $\omega^t = (0.4, 0, 0.6)$  (b) and  $\omega^t = (0.8, 0, 0.2)$  (c).



Fig. 2. The cactus graph used in the proof of the NP-completeness claim.

nonseparable. A block is a maximal nonseparable graph. A cactus graph is a graph in which every block is an edge or a cycle of three or more nodes (see e.g., Brandstadt, Le, & Spinrad, 2000). In Brandstadt et al. (2000) cactus graphs are considered as *almost trees* and classified in the lowest level difficulty class of graphs, just after the acyclic ones. The decision version of the convex OWASTP on cactus graphs can be stated as follows:

*Input*: A cactus graph G = (V, E) with positive integer weights  $(c_e^1, c_e^2)$  assigned to each edge  $e \in E$ . A vector  $w \ge 0$  of rational components  $(w_1, w_2)$ , with  $w_1 > w_2$ , and a rational  $K \ge 0$ . Let  $f_1(T)$  and  $f_2(T)$  be the weights of any spanning tree T of G computed with respect to  $c_e^1$  and  $c_e^2$ , respectively. Let  $\sigma$  be an ordering of  $f_1(T)$  and  $f_2(T)$  such that  $f_{\sigma_1}(T) \ge f_{\sigma_2}(T)$ .

*Output*: Is there a spanning tree *T* of *G* such that, for the ordering  $\sigma$ ,  $w_1 f_{\sigma_1}(T) + w_2 f_{\sigma_2}(T) \le K$ ?

*Claim*: Problem OWASTP is NP-complete on cactus graphs and p = 2.

The reduction is from Partition with Disjoint Pairs (PDP) which is the following problem: Given *n* pairs of integers  $(a_i, b_i)$ , i = 1, ..., n, is there a subset  $S \subset [1, ..., n]$  (a bi-partition) of the set of indexes such that:  $\sum_{i \in S} a_i + \sum_{i \notin S} b_i = \frac{Q}{2}$ , where  $\sum_{i=1}^n (a_i + b_i) = Q$ ? Problem PDP is NP-hard (see e.g., Karp, 1972; Richey, 1990;

Problem PDP is NP-hard (see e.g., Karp, 1972; Richey, 1990; Richey & Punnen, 1992). We observe that PDP is also known as *Alternating Partition* which was proved to be NP-complete in Garey and Johnson (1979).

**Proof.** Given an instance of PDP, construct the (very simple) cactus graph in Fig. 2. Set  $w_1 = 1$ ,  $w_2 = 1 - \delta$ , with  $0 < \delta < 1$  and rational. Let  $K = Q(1 - \frac{\delta}{2})$ .

For each block *i* assign to edge  $(i, j_i)$  weights  $(a_i, b_i)$ ; while to edge  $(i, h_i)$  assign weights  $(b_i, a_i)$ . To all other edges of *G* assign weights (0, 0). Given a solution of PDP, we can construct a solution of OWASTP as follows: if  $i \in S$  the corresponding edge in the spanning tree *T* in block *i* is  $(i, j_i)$ , otherwise add to *T*  $(i, h_i)$ . Then, since *T* must be a spanning tree of *G*, we must add edges  $(j_i, h_i)$  and all the edges (i, 0), i = 1, ..., n. The weight of *T* w.r.t. the first component of the edge weights is  $f_1(T) = \sum_{i \in S} a_i + \sum_{i \notin S} b_i = \frac{Q}{2}$ . Similarly,

$$f_2(T) = \sum_{i \notin S} a_i + \sum_{i \in S} b_i = \frac{Q}{2}.$$
 Hence, for the ordering  $\sigma$  we have:  
$$f_{\sigma_1}(T) + (1 - \delta) f_{\sigma_2}(T) = Q\left(1 - \frac{\delta}{2}\right) = K.$$

Conversely, if we have a solution of OWASTP, *T*, there must exist a subset *S* of the *n* blocks for which edges  $(a_i, b_i)$  belong to *T*,  $i \in S$ ; while for the other blocks (i.e.,  $i \notin S$ ) edges  $(b_i, a_i)$  belong to *T*. Actually, since  $\sum_{i=1}^{n} (a_i + b_i) = Q$ , for any spanning tree *T* such that  $f_1(T) \neq f_2(T)$ , given the ordering  $\sigma$  we have  $f_{\sigma_1}(T) > Q/2$  and  $f_{\sigma_2}(T) < Q/2$ . Suppose that  $f_{\sigma_1}(T) = \frac{Q}{2} + \Gamma$  and  $f_{\sigma_2}(T) = \frac{Q}{2} - \Gamma$ ,  $\Gamma > 0$ . Then, computing the objective function of OWASTP we have:

$$f_{\sigma_1}(T) + (1-\delta)f_{\sigma_2}(T) = \frac{Q}{2} + \Gamma + (1-\delta)\left(\frac{Q}{2} - \Gamma\right)$$
$$= Q\left(1 - \frac{\delta}{2}\right) + \delta\Gamma > K.$$

Since OWASTP belongs to NP, then OWASTP is NP-complete.

We conclude this section by recalling that, according to Yu (1998), if p is unbounded the Min–Max Spanning Tree Problem is strongly NP-hard even for grid graphs (reduction from 3-partition problem which is strongly NP-hard). Following this result, OWASTP is also strongly NP-hard with unbounded number of criteria for grid graphs and this would prevent to find a FPTAS.

## 3. OWASTP formulations

In this section we present several formulations for OWASTP. All of them are MILP formulations, which integrate a mixed integer linear programing STP formulation within a generic mathematical programing formulation for an OWA combinatorial problem (Fernández et al., 2014). We start with the catalogue of STP formulations and then we give the mathematical programing formulations for OWA combinatorial optimization problems that we have used.

## 3.1. Mixed integer linear programing formulations for STP

Many alternative MILP formulations have been proposed for STP. For an overview of the possible alternatives and the properties in each case, the interested reader is addressed to the excellent book chapter by Magnanti and Wolsey (1995) where many of them are presented and compared.

It is well-known that STP formulations exist with the integrality property. Unfortunately, when they are embedded within an OWAP framework the integrality property is lost, so explicit integrality conditions are needed. Alternative STP formulations without such property may now be superior. This explains why some of the formulations we have used lack the integrality property. The criterion that has guided the selection of the formulations is either their good theoretical properties or some characteristic that seemed useful as, for instance, a small number of variables or constraints. We start with two well-known models, the first one derived from the matroid polyhedron (Edmonds, 1970; 1971) and the second one proposed by Martin (1991), both of which having the integrality property. Then we present three existing formulations without the integrality property, based, respectively, on cutset inequalities, flow balance equations and Miller–Tuker–Zemlin inequalities (Miller, Tucker, & Zemlin, 1960). We present another STP formulation based on a relaxation of the formulation proposed by Martin (1991), which uses considerably fewer variables.

All formulations use design variables *x* to represent the edges of the spanning trees. Let  $x_e$ ,  $e \in E$  be a binary variable equal to 1 if edge e = (u, v) is in the spanning tree, and zero otherwise. Some formulations use additional variables related to the arcs of the directed network, D = (V, A) with the same node set as the original undirected *G* and set of arcs *A*, containing two arcs associated with each edge of *E*, i.e.,  $A = \{(u, v), (v, u) \mid (u, v) \in E\}$ .

Throughout we will use the following standard notation. Given a subset of nodes  $S \subset V$ , E(S) and A(S) respectively denote the subsets of edges of E and arcs of A with both end-nodes in S, i.e.,  $E(S) = \{e = (u, v) \in E : u, v \in S\}$  and  $A(S) = \{(u, v) \in A : u, v \in S\}$ . The *cut-set associated with*  $S \subset V$ ,  $\delta(S) = \{e = (u, v) \in E \mid (u \in S, v \in V \setminus S) \text{ or } (v \in S, u \in V \setminus S)\}$ , contains all edges with one node in S and the other node outside S. When working on the directed network D, for  $S \subset V$ , we let  $\delta^+(S) = \{(u, v) \in A \mid u \in S, v \in V \setminus S\}$  denote the cutset directed out of S and  $\delta^-(S) = \{(u, v) \in A \mid u \in V \setminus S\}$  the cutset directed into S. Directed cuts will also be referred to as *dicuts*.

Next we focus on the domains that characterize feasible solutions in each case.

The domain in the subtour elimination formulation is:

$$\mathcal{T}^{sub}: \quad \sum_{e \in E} x_e = n - 1 \tag{1a}$$

$$\sum_{e \in E(S)} x_e \le |S| - 1 \qquad \emptyset \ne S \subset V \tag{1b}$$

$$x_e \ge 0 \qquad e \in E \tag{1c}$$

The cardinality constraint (1a) imposes that exactly n - 1 edges are chosen. Constraints (1b) ensure that the solution contains no cycle. The number of such constraints is exponential on the number of nodes. However, they can be separated in polynomial time by solving a series of minimum (*s*, *t*)-cut problems. An effective algorithm can be implemented using a Gomory–Hu cut tree (Hu, 1974).

It is well-known that all the extreme points in the above domain are integer and that formulation  $T^{sub}$  is stronger than the formulation where inequalities (1b) are replaced by the cut-set constraints  $\Sigma_{e \in \delta(S)} x_e \ge 1$ , that we denote  $T^{cut}$ , which may have fractional extreme points (Magnanti & Wolsey, 1995).

The extended formulation of Martin (1991) models an arborescence rooted at each node  $k \in V$ , in which arcs follow the direction from the leaves to the root. The arcs of such arborescences are then related to the design *x* variables. For  $k \in V$ ,  $(u, v) \in E$ , let  $q_{kuv}$ and  $q_{kvu}$  be decision variables that, respectively, indicate whether or not arcs (u, v) and  $(v, u) \in A$  belong to the arborescence rooted at *k*. The domain of the K. Martin (KM) formulation is the following:

$$T^{km}: \sum_{e \in F} x_e = n - 1$$
(2a)

$$q_{kuv} + q_{kvu} = x_{uv} \qquad k \in V, (u, v) \in E$$
(2b)

$$\sum_{(u,\nu)\in\delta^+(u)}q_{ku\nu}\leq 1 \qquad k\in V, u\in V: u\neq k$$
(2c)

$$\sum_{(k,\nu)\in\delta^+(k)}q_{kk\nu}\leq 0 \qquad k\in V$$
(2d)

$$x_e \ge 0 \qquad e \in E$$
 (2e)

$$q_{ku\nu} \ge 0 \qquad k \in V, \, (u, \nu) \in A \tag{2f}$$

Constraint (2a) ensures that the tree has n - 1 edges. On the other hand, constraints (2b) indicate that the arcs that are used in the arborescences are precisely the ones associated with the n - 1 selected undirected edges. In other words, the underlying undirected graph supporting all the arborescences is exactly the same, so all the arborescences use exactly n - 1 arcs, and the only differences among arborescences are the directions of the arcs, but not the edges on the undirected graph that are used. For each arborescence, (2c) impose that no more than one arc leaves any node different from the root k, while (2d) forbids any arc leaving the root node k. Hence, these constraints imply that for each arborescence, the component containing the root node does not contain any cycle. Since each node is the root of one arborescence, (2b) guarantee that the selected undirected edges contain no cycle and, by (2a), the solutions define spanning trees.

While formulation  $\mathcal{T}^{km}$  has the integrality property, it has an  $O(n^3)$  number of both q variables and constraints (2b). As the size of the graph increases, this number can be prohibitive. When the integrality property is lost because of the addition of new constraints, the computational burden for solving a formulation with such a large number of variables and constraints may become too high.

The Miller-Tucker-Zemlin (MTZ) inequalities are an alternative to the exponential size family of constraints in (1b), to guarantee the connectivity of the solutions and thus prevent cycles. These constraints were initially proposed by Miller et al. (1960) in the context of the Traveling Salesman Problem. They have been adapted to other problems and reinforced by different authors (see e.g. Laporte, 1992; Landete & Marín, 2014). In particular, they have been used by Gouveia (1995) for the Hop-Constrained Spanning Tree Problem, which is a generalization of STP in which the paths starting at a specified root node r are restricted to have no more than p edges. The MTZ formulation for STP builds an arborescence rooted at a specified node  $r \in V$ , in which arcs follow the direction from the root to the leaves. It uses binary variables to represent the arcs of the arborescence. Each edge  $(u, v) \in E$ , is associated with a pair of binary variables,  $y_{uv}$  and  $y_{vu}$ , which take the value 1 if and only if arcs (u, v) and  $(v, u) \in A$  belong to the arborescence, respectively. In addition, it uses continuous variables  $l_u$ , denoting the position that node *u* occupies in the arborescence with respect to r. Since, in principle, there is no pre-specified root node, below r denotes any arbitrarily selected node. The domain of this formulation is given by the following set of constraints:

$$T^{mtz}: \quad \sum_{e \in E} x_e = n - 1 \tag{3a}$$

$$\sum_{\nu,u)\in\delta^{-}(u)}y_{\nu u}=1 \qquad u\in V\setminus\{r\}$$
(3b)

 $y_{uv} + y_{vu} = x_{uv} \qquad (u, v) \in E \tag{3c}$ 

$$l_{\nu} \ge l_{u} + 1 - n(1 - y_{u\nu})$$
  $(u, \nu) \in A$  (3d)

$$u_u = 1$$
  $u = r$  (3e)

$$2 \le l_u \le n \qquad u \in V \setminus \{r\} \tag{3f}$$

$$y_{uv} \in \{0, 1\}$$
  $(u, v) \in A$  (3g)

$$x_e \in \{0,1\} \qquad e \in E \tag{3h}$$

Constraint (3a) ensures that the tree has n-1 edges. Eq. (3b) imposes that each node apart from the root is reached by one single arc, while (3c) guarantee that an edge is selected if any of its two arcs is selected. Constraints (3d) state that if an arc (u, v) is selected the position in the tree of v is higher than the position of u. Finally, (3e) and (3f) assign appropriate bounds to variables  $l_u$ , to ensure that the relative position of the root node in the tree is 1 and that the position of any other node is greater than or equal to 2 and does not exceed the number of nodes.

The flow-based STP formulation we present below (see Magnanti & Wolsey, 1995) is based on the formulation of Gavish (1983) for the capacitated minimal directed tree problem, and was used by Galand and Spanjaard (2012) for OWASTP. In addition to the binary design variables x, the formulation uses continuous flow variables  $\varphi$  defined on the arcs of the directed network D = (V, A). There is a single source node, which is an arbitrarily selected node  $r \in V$ , with inflow n - 1. All other nodes have a demand of one unit. For each  $(u, v) \in A$  the decision variable  $\varphi_{uv}$  represents the amount of flow through arc (u, v). Then the domain of the flow formulation for STP is:

$$T^{flow}: \qquad \sum_{e \in E} x_e = n - 1 \tag{4a}$$

$$\sum_{(r,\nu)\in\delta^+(r)}\varphi_{r\nu}-\sum_{(u,r)\in\delta^-(r)}\varphi_{ur}=n-1$$
(4b)

$$\sum_{(u,v)\in\delta^+(u)}\varphi_{uv} - \sum_{(v,u)\in\delta^-(u)}\varphi_{vu} = -1 \qquad u\in V\setminus\{r\}$$
(4c)

$$\varphi_{uv} \le (n-1)x_{uv} \qquad (u,v) \in E \tag{4d}$$

$$\varphi_{\nu u} \le (n-1)x_{u\nu} \qquad (u,\nu) \in E \tag{4e}$$

$$\varphi_{uv} \ge 0 \qquad (u, v) \in A \tag{4f}$$

$$x_e \in \{0, 1\} \qquad e \in E \tag{4g}$$

Again, constraint (4a) ensures that exactly n - 1 edges are selected. The block of constraints (4b) and (4c) guarantees that n - 1 units of flow leave the source node r and that at least one unit of flow arrives to every other node. The main role of these constraints is to guarantee that the graph induced by the arcs through which the flow circulates is connected and all nodes are "covered". Constraints (4d) and (4e) extend these two properties to the graph induced by the x variables, by imposing that all the edges used for sending flow in some direction are activated.

Concerning domain  $T^{flow}$  note that, because of the flow constraints (4b) and (4c), the removal of constraint (4a) would not change the set of optimal solutions (as opposed to the case of the maximum STP). However, constraint (4a) reinforces considerably the linear relaxation of formulation  $T^{flow}$ , so it is kept in the formulation. Another improvement consists of replacing (4d) and (4e) by the tighter set of constraints:

$$\varphi_{uv} + \varphi_{vu} \le (n-1)x_{uv} \quad (u,v) \in E : u = r \lor v = r$$
(4d')

$$\varphi_{uv} + \varphi_{vu} \le (n-2)x_{uv} \quad (u,v) \in E : u \ne r \land v \ne r$$
(4e')

## 3.1.1. An alternative formulation for STP

Below we present an alternative formulation for STPs, which inherits some of the ideas behind the  $T^{km}$  formulation without requiring  $O(n^3)$  variables. In particular, instead of building one arborescence for each node, we arbitrarily set one single root node

 $r \in V$  and build one single arborescence rooted at r. The arcs of such an arborescence are determined by the subset of variables  $q_{ruv}$ ,  $(u, v) \in A$ . Since r is fixed, in the following we remove the first index and simply denote these variables by  $q_{uv}$ ,  $(u, v) \in A$ . Indeed, equality (2a), plus the subset of constraints (2b), (2c) and (2d) associated with k = r defines a relaxation to formulation  $\mathcal{T}^{km}$ , which uses  $O(n^2)$  variables. Unfortunately, such relaxation is not valid for STPs, as it may produce solutions which are not associated with connected sets of arcs. Luckily, this weakness can be easily overcome by including the following dicut inequalities:

$$\sum_{(u,\nu)\in\delta^+(S)}q_{u\nu}\geq 1, \qquad S\subseteq V\setminus\{r\},$$

which guarantee the connectivity of the obtained solutions (at least one arc will *exit* from any subset of nodes *S* not containing the root node) and thus, the validity of the formulation. The formulation is then as follows:

$$\mathcal{T}^{km2}: \qquad \sum_{(u,v)\in E} x_{uv} = n-1 \tag{6a}$$

$$q_{uv} + q_{vu} = x_{uv} \qquad (u, v) \in E \tag{6b}$$

$$\sum_{u,v)\in\delta^+(u)} q_{uv} \le 1 \qquad u \in V \setminus \{r\}$$
(6c)

$$\sum_{(r,\nu)\in\delta^+(r)}q_{r\nu}\leq 0$$
(6d)

$$\sum_{(u,\nu)\in\delta^+(S)} q_{u\nu} \ge 1 \qquad \emptyset \neq S \subset V \setminus \{r\}$$
(6e)

$$x_{uv} \ge 0 \qquad (u, v) \in E \tag{6f}$$

$$q_{uv} \ge 0 \qquad u, v \in V \tag{6g}$$

#### Remark 1.

- (a) The only difference between formulations  $T^{mtz}$  and  $T^{km2}$  is the way in which subtours are prevented. The former uses the Miller–Tucker–Zemlin inequalities, which are known to be weaker than cut-type constraints used in the latter. This indicates that formulation  $T^{mtz}$  is weaker than  $T^{km2}$ . Below we provide a stronger evidence of the superiority of  $T^{km2}$ over  $T^{mtz}$ , as we will see that  $T^{km2}$  has the integrality property, even if some redundancies are eliminated.
- (b) For any  $u \in V \setminus \{r\}$  the constraint (6e) corresponding to the set  $S = \{u\}$  reduces to  $\sum_{(u,v)\in\delta^+(u)} q_{uv} \ge 1$ . Together with constraints (6c) this implies that  $\sum_{(u,v)\in\delta^+(u)} q_{uv} = 1$  for all  $u \in V \setminus \{r\}$ . Observe, however, that the new set of constraints (6e) together with (6a) and (6b) already imply that  $\sum_{(u,v)\in\delta^+(u)} q_{uv} = 1$  for all  $u \in V \setminus \{r\}$ . To see this, note first that if we add all the constraints (6e) associated with singletons  $S = \{u\}$  with  $u \in V \setminus \{r\}$  we get

$$\sum_{u\in V\setminus\{r\}}\sum_{(u,\nu)\in\delta^+(u)}q_{u\nu}=\sum_{(u,r)\in\delta^-(r)}q_{ur}+\sum_{(u,\nu)\in A(V\setminus\{r\})}q_{u\nu}\geq n-1.$$

Thus, we have,

$$\begin{split} n-1 &\leq \sum_{(u,r)\in\delta^{-}(r)} q_{ur} + \sum_{(u,\nu)\in A(V\setminus\{r\})} q_{u\nu} \\ &\leq \sum_{(r,\nu)\in\delta^{+}(r)} q_{r\nu} + \sum_{(u,r)\in\delta^{-}(r)} q_{ur} + \sum_{(u,\nu)\in A(V\setminus\{r\})} q_{u\nu} \\ &= \sum_{(u,\nu)\in A} q_{u\nu} = \sum_{(u,\nu)\in E} x_{u\nu} = n-1, \end{split}$$

where the last two equalities follow from constraints (6b) and (6a), respectively.

Hence, we can conclude that  $\sum_{(u,v)\in\delta^+(r)}q_{uv}=0$  and  $\sum_{(u,v)\in\delta^+(u)}q_{uv}=1$  for all  $u \in V \setminus \{r\}$ , since otherwise we would reach a contradiction.

The above remark indicates that the dicut constraints (6e) make the sets of constraints (6c) and (6d) unnecessary. Hence, STP formulation which emanates from the above discussion is:

$$T^{dc}: \qquad \sum_{(u,v)\in E} x_{uv} = n-1 \tag{7a}$$

$$q_{uv} + q_{vu} = x_{uv} \qquad (u, v) \in E \tag{7b}$$

$$\sum_{(u,v)\in\delta^+(S)} q_{uv} \ge 1 \qquad \emptyset \neq S \subset V \setminus \{r\}$$
(7c)

$$x_{uv} \ge 0 \qquad (u, v) \in E \tag{7d}$$

$$q_{uv} \ge 0 \qquad u, v \in V \tag{7e}$$

The reader may observe that formulation  $T^{dc}$  can be readily transformed into the *directed cut* formulation of Magnanti and Wolsey (1995) by just changing the directions of the arcs of the arborescence and, thus, directing the arcs from the root r to the leaves, instead of from the leaves to the root. Since the directed cut formulation of Magnanti and Wolsey (1995) has the integrality property, so does formulation  $T^{dc}$ . In its turn, this implies the integrality of the domain of  $T^{km2}$ .

The number of dicut constraints (7c) is exponential on |V|. Nevertheless, they can be incorporated into the formulation only if needed via an efficient separation oracle, as they can be separated in polynomial time by finding the Gomory–Hu cut tree (Hu, 1974).

## 3.1.2. Comparison of formulations

Let  $P(\mathcal{T}^{(\cdot)})$  denote the polyhedron associated with the linear programing relaxation of formulation  $\mathcal{T}^{(\cdot)}$ . Except for formulation  $\mathcal{T}^{sub}$ , all other formulations above are extended formulations, in the sense that, besides the design *x* variables, additional sets of variables are used. For comparing all the formulations in the same space we project the polyhedra associated with the extended formulations onto the space of the *x* variables, and denote by  $P_X(\mathcal{T}^{(\cdot)})$  the projected polyhedron associated with formulation  $\mathcal{T}^{(\cdot)}$ .

Several of the formulations described above have the integrality property, namely formulations  $T^{sub}$ ,  $T^{km}$  and  $T^{km2}$ . This means that  $P_x(T^{sub}) = P_x(T^{km}) = P_x(T^{km2})$ . In its turn, each of these formulations is tighter than any of the formulations without integrality property. That is,  $P_x(T^{km2}) \subset P_x(T^{mtz})$  and  $P_x(T^{km2}) \subset$  $P_x(T^{flow})$ . Below we compare  $P_x(T^{mtz})$  and  $P_x(T^{flow})$ , as we have not seen such comparison in the literature.

The example of Fig. 3 illustrates that  $P_x(T^{flow}) \not\subseteq P_x(T^{mtz})$ . The components of a *x* vector such that  $\sum_{e \in E} x_e = n - 1$  are given next to each edge. Taking r = 5 as the root node, the flow  $\varphi_{53} = \varphi_{54} = 2$ ,  $\varphi_{31} = \varphi_{42} = 1$ , together with *x*, define a feasible solution to formulation  $P_x(T^{flow})$ . However, there is no feasible *y* vector that together with the depicted *x* vector satisfies constraints (3b) and (3c).



**Fig. 3.** Fractional *x* solution with  $\sum_{e \in E} x_e = n - 1$ .



**Fig. 4.** Solution *x* of  $T^{mtz}$  formulation in the complete graph (n = 5) of the above example.

On the other hand, the example depicted in Fig. 4 shows that  $P_X(T^{mtz}) \not\subseteq P_X(T^{flow})$ , i.e. the two formulations are not related in that there exist feasible solutions to  $P_X(T^{flow})$  that do not give rise to feasible solutions to  $P_X(T^{mtz})$  and the other way around.

Consider a complete graph with n = 5 nodes and cost matrix:

	1	0	31	19	33	67	
	1	31	0	57	40	38	
<i>C</i> =		19	57	0	2	18	
		33	40	2	0	13	
		67	38	18	13	0	)

The optimal solution to the linear relaxation of  $T^{mtz}$  is given by:  $x_{12} = 1$ ,  $x_{34} = 1$ ,  $x_{35} = 1$ ,  $x_{45} = 1$ ;  $y_{12} = 1$ ,  $y_{34} = 0.5$ ,  $y_{35} = 0.5$ ,  $y_{43} = 0.5$ ,  $y_{45} = 0.5$ ,  $y_{53} = 0.5$ ,  $y_{54} = 0.5$ ; and  $\ell_1 = 1$ ,  $\ell_2 = 2$ ,  $\ell_3 = 2$ ,  $\ell_4 = 2$ ,  $\ell_5 = 3.5$ .

It is clear that the above solution to  $T^{mtz}$  does not induce a feasible solution to  $T^{flow}$  since the vector x does not produce a connected solution in the graph. Thus we have the following result:

## **Corollary 1.**

$$P_{x}(T^{sub}) = P_{x}(T^{km}) = P_{x}(T^{km2}) \subseteq \begin{cases} P_{x}(T^{mt2}) \\ \neq \\ P_{x}(T^{flow}) \end{cases}$$

## 3.2. Mixed integer linear programing formulations for OWAP

This section presents the OWA formulation that we use for OWASTP. The choice is based on our preliminary experiments for STP and on previous results of Fernández et al. (2014), that show that this formulation outperforms other alternatives when the embedded combinatorial object is the shortest path or the perfect matching problem. In the formulations below we assume that we use the same polyhedron to represent the combinatorial object T, namely the set of spanning trees. This is expressed as  $x \in T$ .

Consider the following binary variables that define the specific positions in the ordering of the sorted cost function values:

 $z_{ij} = \begin{cases} 1 & \text{if cost function ioccupies position } j, \\ 0 & \text{otherwise} \end{cases}$ 

For each  $j \in P$ , let also  $\theta_j$  be a variable representing the value of the objective function sorted at position *j*. Then, OWAP can be formulated as:

$$F^{\theta}: \quad V = \min \sum_{j \in P} \omega_j \theta_j$$
 (8a)

s.t. 
$$\sum_{i\in P} Z_{ij} = 1$$
  $j \in P$  (8b)

$$\sum_{j\in P} z_{ij} = 1 \qquad i \in P \tag{8c}$$

$$C^{i}x \le \theta_{j} + M(1 - \sum_{k \ge j} z_{ik}) \qquad i, j \in P$$
(8d)

$$\theta_j \ge \theta_{j+1} \qquad j \in P : j 
(8e)$$

$$x \in T$$
 (8f)

$$\theta_j \ge 0 \qquad j \in P \tag{8g}$$

$$z \in \{0, 1\}^{p \times p} \tag{8h}$$

The objective function (8a) minimizes the weighted average of sorted objective function values, provided that  $\theta_j$ ,  $j \in P$ , are enforced to take on the appropriate values. Constraints (8b) and (8c) define a permutation of the cost functions, by placing one single cost function at each position and each cost function at one single position of the sequence. Constraints (8d) relate cost function values with the values placed in the sorted sequence. Constraints (8e) are optimality cuts which help the resolution of  $F^{\theta}$ , as explained in Fernández et al. (2014).

For comparison purposes in our computational experiments, below we present the formulation used by Galand and Spanjaard (2012) for OWASTP. This formulation uses the above *z* binary variables plus an additional set of continuous variables  $y = (y_{ij})_{i,j\in P} \in \mathbb{R}^{p \times p}$ , where  $y_{ij}$  denotes the value of cost function *i* if it occupies the *j*th position in the ordering. The formulation is as follows:

$$F^{GS}: \qquad V = \min \sum_{j \in P} \omega_j \sum_{i \in P} y_{ij}$$
(9a)

s.t. 
$$\sum_{i\in P} z_{ij} = 1$$
  $j \in P$  (9b)

$$\sum_{j \in P} Z_{ij} = 1 \qquad i \in P \tag{9c}$$

$$\sum_{i \in P} y_{ij} \ge \sum_{i \in P} y_{ij+1} \qquad j \in P : j 
(9d)$$

$$y_{ij} \le M z_{ij}$$
  $i, j \in P$  (9e)

$$\sum_{j\in P} y_{ij} = C^i x \qquad i \in P \tag{9f}$$

$$x \in T$$
 (9g)

$$y_{ij} \ge 0 \qquad i, j \in P \tag{9h}$$

$$z \in \{0, 1\}^{p \times p} \tag{9i}$$

The objective function (9a) minimizes the weighted average of sorted objective function values. Constraints (9b) and (9c) are a copy of constraints (8b) and (8c) respectively, and thus define a cost function permutation. Constraints (9d) impose that the sorted values are ordered non-increasingly. Constraints (9e) relate cost function values with the values placed in the sorted sequence. Constraints (9f) ensure that one of the  $y_{ij}$  variables gives precisely the value of the objective function *i*.

Note that the relationship between  $\theta$  in formulation  $F^{\theta}$  and the *y* variables in  $F^{CS}$  is:

$$\theta_j = \sum_{i \in P} y_{ij} \qquad j \in P : j > 1.$$
(10)

The reader should observe that  $F^{\theta}$  and  $F^{GS}$  differ in the way we represent the OWA operator, although as mentioned at the beginning of this section the combinatorial object T is represented by means of the same polyhedron.

Next, we prove two results concerning formulations  $F^{\theta}$  and  $F^{GS}$ . Let us denote by  $\Omega^{\theta}$  and  $\Omega^{GS}$  the domains defined by their respective sets of constraints. We first prove that  $F^{\theta}$  and  $F^{GS}$  have the same set of optimal solutions although  $\Omega^{GS} \subset \Omega^{\theta}$ . This property no longer holds for the respective relaxations, where everything remains unchanged except for the *z* variables, which are allowed to take continuous values, i.e.  $z_{ij} \geq 0$ ,  $i, j \in P$ . In particular, we will see that  $\Omega_{LR}^{GS} \subset \Omega_{LR}^{\theta}$ , where  $\Omega_{LR}^{\theta}$  and  $\Omega_{LR}^{GS}$  denote the respective continuous relaxed domains. Moreover, in general, the sets of optimal solutions of the linear relaxations for the objective functions (8a) and (9a) do not coincide.

**Property 2.** Every optimal solution to  $F^{CS}$  is also optimal to  $F^{\theta}$  and conversely.

**Proof.** Given the relationship (10) between  $\theta$  and y variables, in  $\Omega^{\theta}$  we can substitute Constraints (8d) by  $C^{i}x \leq \sum_{i \in P} y_{ij} + M(1 - \sum_{k \geq j} z_{ik})$ ,  $i, j \in P$ .

• We prove first that  $\Omega^{GS} \subseteq \Omega^{\theta}$ , that is, every solution  $(x, z, y) \in \Omega^{GS}$  (not necessarily optimal) is such that  $(x, z, y) \in \Omega^{\theta}$ . Observe that it suffices to prove that every  $(x, z, y) \in \Omega^{GS}$ , with y and  $\theta$  related by (10), satisfies

$$C^{i}x \leq \sum_{i \in P} y_{ij} + M(1 - \sum_{k \geq j} z_{ik}) \qquad i, j \in P$$
(9d').

Let  $\hat{x}$  be a feasible solution in  $\mathcal{T}$  and  $\hat{z}$  a permutation that sorts the cost function values of x. Then, for fixed  $\hat{x}$  and  $\hat{z}$  values there is a unique  $\hat{y}$  since, according to (9e) and (9f) there is at most one  $j \in P$  such that  $y_{ij} \neq 0$  for each  $i \in P$ . It follows that such  $(\hat{x}, \hat{z}, \hat{y})$  verifies (9d').

• Next we prove that every optimal solution of  $F^{\theta}$ , satisfies that  $(x, z, y) \in \Omega^{GS}$ , after performing the change of variable given by (10). For this, it is sufficient to prove that every optimal solution  $(x, z, y) \in \Omega^{\theta}$  verifies (9e) and (9f). Let  $\hat{x}$  be a feasible solution in T. Then, there exists a permutation  $\hat{z}$  that sorts the cost function values of  $\hat{x}$  in non increasing order. Now, we give values to the  $\hat{\theta}$  variables according to this ordered sequence, and we determine the  $\hat{y}$  values by means of  $\hat{y}_{ij} = \hat{\theta}_j \hat{z}_{ij}$   $i, j \in P$ . From here, it follows that  $(\hat{x}, \hat{z}, \hat{y})$  verifies (9d)–(9f). In addition, we note that, in general, for fixed  $\hat{x}$  and  $\hat{z}$ , the polyhedron given by (8d) and (8e) is unbounded and thus  $\Omega^{\theta} \notin \Omega^{GS}$ .

## **Property 3.** $\Omega_{LR}^{GS} \subset \Omega_{LR}^{\theta}$ .

**Proof.** First, we observe that  $\Omega_{LR}^{\theta} \not\subseteq \Omega_{LR}^{GS}$  since, otherwise, the optimal solution of  $F_{LR}^{\theta}$  for the graph in Example 1 (with value 8.6 when  $\omega = (0.8, 0, 0.2)$ ) could not have a smaller value than the optimal solution of  $F_{LR}^{GS}$  (with value 9.4).

Next, we prove that every feasible solution  $(x, z, y) \in \Omega_{LR}^{GS}$  is such that  $(x, z, y) \in \Omega_{LR}^{\theta}$ , once the change of variable given by (10) is done.

Indeed, we have to prove that any  $(x, z, y) \in \Omega_{IR}^{GS}$  verifies

$$C^{i}x \leq \sum_{i \in P} y_{ij} + M\left(1 - \sum_{k \geq j} z_{ik}\right) \qquad i, j \in P$$
(9d').

Let  $\hat{x}$  be a feasible solution in  $\mathcal{T}$  and  $\hat{z}$  a fractional vector. Since  $C^i x = \sum_{i \in P} y_{ij}$  and  $(1 - \sum_{k \geq j} z_{ik})$  is greater than or equal to zero, it is clear that constraint (9d') is verified.

The above result proves that the linear relaxation of  $\Omega^{GS}$  is stronger than that of  $\Omega^{\theta}$ , although the two formulations share the same optimal integer values. Nevertheless, as we shall show in the computational experiments, formulation  $F^{\theta}$  provides much better results in terms of running times and number of optimal solutions found. The reason may be the smaller number of variables used in the second formulation.

To conclude this section we state the relationships between the different formulations that derived from the combination of some OWA representation and any of STP polytopes described above. To this end, let us denote by  $P_{XZ}(\Omega^{(\cdot)})$  the projection onto the space of the *x*, *z* variables of the linear relaxation of an OWA polytope built on the corresponding  $T^{(\cdot)}$  polytope for STP. The following property states the relationships among them.

#### Property 4.

$$P_{xz}(\Omega^{sub}) = P_{xz}(\Omega^{km}) = P_{xz}(\Omega^{km2}) \subseteq \begin{cases} P_{xz}(\Omega^{flow}) \\ \neq \\ P_{xz}(\Omega^{mtz}) \end{cases}$$
(11)

## 3.2.1. Enhancements and valid inequalities for OWAP

Formulation  $F^{\theta}$  admits other enhancements like removing some redundant variables, adding valid inequalities, etc. First, we observe that since system (8b) and (8c) contains exactly 2p - 1 linearly independent equations, the above permutation can also be represented without variables  $z_{i1}$ , for all  $i \in P$ , which can be replaced by  $1 - \sum_{j \in P: j > 1} z_{ij}$ . In this way, system (8b) and (8c) can also be rewritten as

$$\sum_{i\in P} z_{ij} = 1 \qquad j \in P : j > 1, \tag{12}$$

$$\sum_{i\in P} z_{ij} \le 1 \qquad i \in P.$$
(13)

Second, constraints (8c) and (8e) can be removed from  $F^{\theta}$  without changing the set of optimal solutions. We denote by  $F_{R1}^{\theta}$  formulation  $F^{\theta} \setminus \{(8e)\}$  and by  $F_{R2}^{\theta}$  formulation  $F^{\theta} \setminus \{(8e)\}$ . Note that formulations  $F_{R1}^{\theta}$  and  $F_{R2}^{\theta}$  admit some solutions that are infeasible in  $F^{\theta}$  (e.g. a solution where  $\theta_j \leq \theta_{j+1}$  for some *j*). However, these two new formulations have fewer constraints and could be more efficient in a branch-and-bound algorithm.

Finally, we present some valid inequalities that can be added to the above OWAP in order to improve the bound of the linear relaxation and/or to reduce the search space in the branch-andbound tree.

• Constraints related to bounds of cost function values. Let  $l_i(u_i)$  denote the minimum (maximum) objective value relative to cost function  $i \in P$ , respectively. It is clear that  $l_i(u_i)$  are valid lower

(upper) bounds on the value of objective *i*, independently of the position of cost function *i* in the ordering. Therefore we obtain the following two sets of constraints which are valid for OWAP:

$$l_i \le C^i x \le u_i \qquad i \in P \tag{14}$$

• Constraints related to bounds of values in specific positions. Let  $l_j^{\pi}$   $(u_j^{\pi})$  denote the *j*th lowest (largest) value of  $l_i$   $(u_i)$ . Then,  $l_j^{\pi}$   $(u_j^{\pi})$  is a valid lower (upper) bound of the objective function sorted in position *j*, that is

$$l_i^{\pi} \le \theta_i \le u_i^{\pi} \qquad j \in P \tag{15}$$

• There are also different bounds on the value of the cost function *i* and the value of the cost function sorted in position *j*:

$$\sum_{j\in P} \max\{l_i, l_j^{\pi}\} z_{ij} \le C^i x \le \sum_{j\in P} \min\{u_i, u_j^{\pi}\} z_{ij} \qquad i \in P$$
(16)

#### 4. Computational experience

Next, we report on the results of some computational experiments we have run, in order to compare empirically the proposed formulations and reinforcements. We have studied OWASTP combining the different formulations proposed for STP and OWAP. First of all we have chosen the best formulation, according to Fernández et al. (2014), among those proposed for OWAP, namely  $F_{R2}^{\theta}$ . We recall that the goal of this paper is the analysis of some STP formulations within the OWAP framework.

In our computational experience we study three particular cases of the OWA operator already considered in multiobjective optimization (see e.g., Galand & Spanjaard, 2012). We study first the k-centrum criterion (Tamir, 2000) that evaluates the sum of the k greatest objective functions. This criterion is monotonic and convex since the sorting weights  $(w_1 = \cdots = w_k = 1/k, w_{k+1} = \cdots =$  $w_p = 0$ ) are in decreasing order. For that reason this operator can be modeled using the linearization of the OWA function with monotonic weights given by Ogryczak and Sliwinski (2003) and Ogryczak and Tamir (2003), which avoids the use of binary variables. Then, we study two non-monotonic and non-convex criteria namely the k-trimmed mean (Galand & Spanjaard, 2012) defined as  $\sum_{i=k+1}^{p-k} (p-2k)^{-1} y_i$  and the Hurwicz criterion (Hurwicz, 1951), defined as  $\alpha \max_{i \in P} y_i + (1 - \alpha) \min_{i \in P} y_i$ . These criteria are of special interest for being non-convex since the sorting weights ( $w_1 =$  $\alpha, w_2 = \cdots = w_{p-1} = 0, w_p = 1 - \alpha \text{ and } w_1 = \cdots = w_k = 0, w_{k+1} = 0$  $\dots = w_{p-k-1} = 1/(p-2k), \ w_{p-k} = \dots = w_p = 0, \ \text{respectively}) \ \text{are}$ not in non-increasing order (Grzybowski et al., 2011, Puerto & Tamir, 2005).

For our computational experiments we have followed the design of Galand and Spanjaard (2012). Thus, the number of objectives ranges in  $|P| \in \{5, 8, 10\}$  and the considered values of k,  $\alpha$  are

- *k*-centrum: |P| = 5 and  $k \in \{1, 3, 4\}$ , |P| = 8 and  $k \in \{2, 4, 7\}$ , |P| = 10 and  $k \in \{3, 5, 8\}$ ,
- Hurwicz:  $|P| \in \{5, 8, 10\}$  and  $\alpha \in \{0.4, 0.6, 0.8\}$ ,
- *k*-trimmed: |P| = 5 and  $k \in \{1, 2\}, |P| = 8$  and  $k \in \{2, 3\}$ .

Graphs are complete with  $|V| \in \{20, 30, 40, 50, 60, 70, 80, 100\}$ and the components of the cost vectors randomly drawn from a uniform distribution on [1, 100]. In addition, for each selection of the parameters (|V|, p), 10 instances were randomly generated. All instances were solved with the MIP Xpress 7.7 optimizer, under a Windows 7 environment in an Intel(R) Core(TM)i7 CPU 2.93 gigahertz processor and 16 gigabytes RAM. Default values were initially used for all parameters of Xpress solver and a time limit of 3600 seconds was set. We have also tested different combinations of parameters for the solver cut strategy and intensity of heuristics but, unless it is specified, the best results were obtained with the parameters of the solver set to the default values.

Throughout the section  $F^{GS}$  denotes the formulation of Galand and Spanjaard (2012) for OWASTP. Otherwise, we denote by  $F^{(.)}$  the combination of the OWA  $F_{R2}^{\theta}$  formulation together with a STP  $T^{(.)}$ . We report results of formulations  $F^{GS}$ ,  $F^{km}$ ,  $F^{cut}$ ,  $F^{mtz}$ ,  $F^{flow}$ , and  $F^{km2}$ .

The separation of the cutset inequalities in formulation  $F^{cut}$  was implemented using a max-flow based algorithm (Gusfield, 1990). Heuristics in Xpress solver were configured with intensity 2 (out of 3) and an initial solution was given to the problem. The initial solution was the minimum cost spanning tree obtained using as edge costs the average costs among all objectives.

We have summarized the results in five sets of three tables each. In each set, the first table refers to OWA 1 (k-centrum), the second one to OWA 2 (Hurwicz) and the third one to OWA 3 (trimmed mean), respectively.

Table 1.c ,  $c \in \{1, 2, 3\}$  show results with OWA c for formulations  $F^{GS}$ ,  $F^{km}$ ,  $F^{cut}$ ,  $F^{mtz}$ ,  $F^{flow}$ , and  $F^{km2}$ . Table 2.c,  $c \in \{1, 2, 3\}$  give results of the best strengthening for  $F^{km2}$ , which consist of inequalities (14), (15) and (16). Analogously, Table 3.c,  $c \in \{1, 2, 3\}$  report the same information but referred to  $F^{mtz}$ . Table 4.c,  $c \in \{1, 2, 3\}$ , display a comparison of our two best formulations with the results reported in Galand and Spanjaard (2012). The results of a last series of experiments with larger graphs with up to |V| = 100 nodes and with up to |P| = 10 objectives are presented in Tables 5.c,  $c \in \{1, 2\}$ . We do not give results of OWA 3 (*k*-trimmed) since Table 4.3 indicates that the instance sizes limit for this criterion is already reached for |P| = 8 objectives.

In order to facilitate the comparison among all tables, best results in each table are marked in bold. In all tables each row summarizes the results corresponding to a group of instances with the same parameters (|P|, |V|,  $\alpha$ ). Columns are grouped in blocks. The first block contains three columns with the values of the instances parameters. In Tables 1–3.c,  $c \in \{1, 2, 3\}$  rows correspond to groups of 10 instances. The first block of columns is followed by a block of four columns for each tested formulation. The columns of each such block are the following. Columns gapR give the percentage relative gap at the root node, computed as  $100(z^* - z_R)/z_R$ , where  $z^*$  denotes the value of the best solution found and  $z_R$  the optimal value of the linear relaxation at the root node. Columns t/gap(#)report average computing times in seconds over the 10 instances of the row (denoted by t). When t is smaller than 10 seconds we report one additional precision digit. If at least one instance reaches the time limit, the number of instances in the group solved to optimality within the time limit is given in brackets (#). In such a case, *t* is computed using the time limit for all unsolved instances. If no instance was solved to optimality, instead of t we report the average optimality gap relative to the lower bound at termination over all the instances of the group, (denoted by gap). Columns  $t^*/gap^*$  show the maximum computing time in seconds over the 10 instances of the row  $(t^*)$ . If at least one instance reached the time limit, instead of  $t^*$  we report the maximum optimality gap, over all the instances of the group (denoted by gap<sup>\*</sup>). Finally, columns *nod* indicate the average number of nodes explored in the branch-and-bound tree in compact format denoting  $a^*10^n$  as aen. The caption just below each block gives the formulation the block refers to.

Table 4.c,  $c \in \{1, 2, 3\}$ , display the results obtained and reported in Galand and Spanjaard (2012) for OWASTP and our two best formulations. Results provided by Galand and Spanjaard (2012) correspond to minimum, average and maximum running times for groups of 30 instances with the same parameters (|P|, |V|). An entry with the symbol "–" indicates that the average execution time was beyond 15 minutes (900 seconds). Block  $F^{GS}$  shows the results reported by Galand and Spanjaard (2012) in IBM ILOG CPLEX 12 without any preprocessing whereas, blocks  $F_{P1}^{CS}$  and  $F_{P2}^{CS}$  give the running times after applying two different preprocessings (shavings) described in that paper. In that case, all the algorithms were implemented in C++ and were run on either an Intel Xeon 2.5 gigahertz personal computer with a memory of 4 gigabytes for five objectives, or an Intel Core 3.0 gigahertz personal computer with a memory of 8 gigabytes for eight objectives. For the sake of comparison and according with https://www.spec.org/cgi-bin/osgresults , the performance indices of the Intel Core 17 3 gigahertz, used in Galand and Spanjaard (2012) and the Intel(R) Core(TM)i7 CPU 2.93 gigahertz, used in our computational results, are rather similar; whereas the performance of the Intel Xeon 2.5 gigahertz is slightly inferior.

The meaning of the columns of Table 5.c,  $c \in \{1, 2\}$  is like in the first three sets of tables.

All data instances and disaggregated results of all experiments are available via e-mail upon request to the authors.

#### 4.1. Tests with the k-centrum criterion

In Table 1.1, the results of block  $F^{GS}$  exhibit small values of gapR for all instance sizes. However, many of these instances remain unsolved after 1 hour of CPU time leaving integrality gaps around 1%. The results of block  $F^{km}$  show that, according to the low average number of explored nodes in the B&B tree, solving the LP relaxation of the problem becomes quite hard so the corresponding gaps at termination remain quite large in comparison with  $F^{GS}$ . We recall that  $F^{km}$  uses  $O(n^3)$  variables and constraints, which can be too high in large graphs. The results of block F<sup>cut</sup> indicate that, in general, this formulation is outperformed by all formulations. Block *F<sup>flow</sup>* behaves similarly to *F<sup>GS</sup>* which suggests that flow MST formulations embedded within an OWA framework for the k-centrum criterion does not exhibit a good performance. On the contrary, block  $F^{mtz}$  shows the best performance in terms of number of instances solved to optimality, running times and optimality gaps. Block *F*<sup>km2</sup> shows a good performance but not as good as F<sup>mtz</sup>.

Next we report on the three most promising strengthening found, which consist on adding valid inequalities (14), (15) to formulations  $F^{km2}$  and  $F^{mtz}$ . Note that reinforcement (16) cannot be applied to the *k*-centrum criteria since we are using here the formulation of Ogryczak and Tamir (2003) that does not require binary variables  $z_{ij}$ . Table 2.1 shows that the performance of  $F^{km2}$  is slightly improved when constraints (15) are added. On the contrary, Table 3.1 shows that the performance of  $F^{mtz}$  is not improved in general adding any of these valid inequalities.

Fig. 5 summarizes the main results of Tables 1.1, 2.1 and 3.1 as follows. To better illustrate the results we have shown in a single figure the information relative to computing time and optimality gap at termination. For this, we have defined the Time/Gap (TG) *performance index* of an instance as  $0.5(\frac{time}{3600} + \frac{gap}{100})$ . This index is represented in a plot where the horizontal axis depicts the size of the instances, namely (|P|, |V|), and the vertical axis corresponds to the scaled values of the index. The reader should observe that whenever for a given instance the gap at termination is null this index reflects its computing time and is always represented below the horizontal line (3600, 0%) because the corresponding instance must have been solved within the time limit. Analogously, a point above the horizontal line (3600, 0%), indicates an instance with a strictly positive gap at the time limit. The average TG performance index of a set of instances is the average of their TG performance values and the worst TG index is the maximum TG value among the instances in the set. For each instance size (|P|, |V|), Fig. 5(a) and (b) plot the average and worst TG performance index, respecTable 1

OWASTP results for the different formulations.

Table 11	Results	for	the	k-centrum	criterion
Tuble 1.1.	nesuits	101	the	R centrum	criterion.

P	V	k	gapR	<i>t/gap</i> (#)	t*/gap*	nod	gapR	<i>t/gap</i> (#)	t*/gap*	nod	gapR	t/gap(#)	t*/gap*(%)	nod	gapR	t/gap(#)	t*/gap*	nod	gapR	t/gap(#)	t*/gap*	nod	gapR	<i>t/gap</i> (#)	t*/gap*	nod
5	40	1	3.0	429	2510	3e5	3.6	184	404	3e3	18.2	3449(1)	33.4	5e6	3.0	479(9)	0.1%	3e5	0.7	3.0	7.2	3e3	0.7	2.1	6.2	5e3
5	40	3	2.7	192	1034	1e5	0.3	148	456	4e2	10.0	3345(1)	26.7	3e6	2.7	137	769	1e5	0.4	1.7	2.8	5e2	0.4	1.4	6.4	2e3
5	40	4	2.3	8.6	56	8e3	0.1	42	384	15	2.8	3241(1)	6.1	3e6	2.3	8.8	65	7e3	0.2	0.8	2.0	8	0.2	0.5	1.7	2e2
5	50	1	2.8	1200(8)	0.4%	3e5	3.2	1224	2171	5e3	29.2	28.9%(0)	32.8	3e6	2.8	429	1624	3e5	0.5	22	153	2e4	0.6	417(9)	0.5%	6e5
5	50	3	2.6	262	1658	6e4	6.0	819(9)	27.6%	6e2	15.5	15.2%(0)	31.6	3e6	2.6	72	268	5e4	0.4	4.4	16	6e3	0.5	173	1636	3e5
5	50	4	2.4	146	1345	6e4	0.1	165	1591	16	3.8	3291(1)	6.8	2e6	2.4	20	137	2e4	0.2	1.5	5.6	7e2	0.2	3.2	19	2e3
5	60	1	3.0	3386(1)	1.5%	5e5	23.3	2939(2)	32.0%	2e3	29.3	29.2%(0)	33.2	2e6	2.9	2502(4)	1.1%	1e6	0.5	75	537	8e4	0.6	694(9)	0.2%	7e5
5	60	3	2.7	2331(5)	0.8%	3e5	19.9	2532(3)	30.5%	2e2	24.9	24.7%(0)	33.0	2e6	2.7	1857(6)	0.7%	6e5	0.4	39	179	3e4	0.4	295	2536	3e5
5	60	4	2.4	744(9)	0.5%	2e5	3.2	369(9)	31.7%	4	3.9	3.5%(0)	6.3	2e6	2.4	430(9)	0.2%	3e5	0.2	1.3	3.0	3e2	0.3	8.3	57	5e3
8	40	2	2.4	3085(3)	1.4%	1e6	5.2	2297(6)	0.5%	3e4	16.0	15.5%(0)	24.8	4e6	2.3	3256(3)	0.9%	2e6	0.9	146	372	2e5	0.9	182	639	3e5
8	40	4	2.1	1230(8)	0.5%	5e5	0.6	956	3125	2e4	11.0	10.5%(0)	20.9	5e6	2.1	1051(8)	0.7%	6e5	0.7	25	126	4e4	0.7	47	256	6e4
8	40	7	1.4	12	77	7e3	0.0	6.0	13	8	2.0	3437(1)	4.2	2e6	1.4	7.4	36	6e3	0.1	1.2	3.1	3e2	0.1	1.1	4.6	5e2
8	50	2	2.1	3424(1)	1.3%	5e5	5.4	3043(2)	1.4%	6e3	17.5	17.2%(0)	27.7	9e6	2.0	3293(1)	0.7%	9e5	0.7	849(9)	0.3%	6e5	0.8	1260(8)	0.9%	1e6
8	50	4	1.8	2754(4)	0.9%	5e5	5.2	2085(9)	0.1%	2e4	13.8	13.4%(0)	25.7	9e6	1.8	2185(5)	0.6%	7e5	0.5	79	503	8e4	0.7	839(8)	0.8%	7e5
8	50	7	1.4	460(9)	0.0%	1e5	0.1	207	1949	62	2.4	2.0%(0)	4.7	8e6	1.4	102	718	7e4	0.1	3.3	18	2e3	0.2	53	387	6e4
8	60	2	2.4	0.8%(0)	2.1%	3e5	25.3	3490(1)	29.8%	0	24.9	24.7%(0)	29.2	5e6	2.1	0.5%(0)	1.1%	9e5	0.6	<b>1832</b> (6)	1.1%	8e5	0.6	1839( <b>7</b> )	0.7%	1e6
8	60	4	2.1	0.5%(0)	1.2%	3e5	11.7	11.6%(0)	26.0%	8e3	21.2	21.1%(0)	28.3	5e6	2.0	3442(1)	0.9%	8e5	0.5	1289( <b>7</b> )	0.7%	5e5	0.5	1159(7)	0.7%	7e5
8	60	7	1.6	330	2388	7e4	3.0	375(9)	29.7%	17	2.8	2.5%(0)	5.4	5e6	1.6	114	842	4e4	0.1	5.4	24	4e3	0.1	192	1744	1e5
				$F^{G}$	S			F <sup>ki</sup>	m			F <sup>ct</sup>	ıt			Fflow	/			$F^n$	ıtz			$F^k$	m2	
Tabl	e 1.2	. Res	ults for t	he Hurwic	z criterion.																					
P	V	α	gapR	t/gap(#)	t*/gap*(%)	nod	gapR	t/gap(#)	t*/gap*(%)	nod	gapR	t/gap(#)	t*/gap*	nod	gapR	t/gap(#)	t*/gap*	nod	gapR	t/gap(#)	t*/gap*	nod	gapR	t/gap(#)	t*/gap*	nod
5	40	0.4	41.5	3303(1)	14.7	5e6	47.5	3102(2)	48.9	1e4	48.5	350	1629	3e3	48.5	108	707	1e5	47.3	16	66	1e3	47.3	21	114	3e3
5	40	0.6	26.9	2432(4)	13.5	4e6	36.0	2366(4)	36.1	1e4	37.4	140	491	1e3	37.3	19	46	3e4	35.9	17	65	1e3	35.9	13	40	1e3
5	40	0.8	8.2	2301(4)	3.6	2e6	20.3	1481(7)	19.7	5e3	22.2	685	2807	6e3	22.2	83	371	8e4	20.4	16	65	2e3	20.4	22	47	3e3
5	50	0.4	40.4	3207(2)	19.4	3e6	48.0	3113(2)	49.6	5e3	48.8	429	945	2e3	48.8	239	816	2e5	47.6	33	70	2e3	47.6	49	163	3e3
5	50	0.6	27.2	3276(1)	13.3	4e6	36.5	3337(1)	37.8	5e3	37.5	820(9)	0.1%	3e3	37.5	772(9)	0.1%	6e5	36.0	34	73	6e3	36.1	48	191	3e3
5	50	0.8	7.4	3310(2)	1.6	2e6	20.3	2542(5)	20.0	4e3	22.2	1661(8)	0.4%	6e3	22.1	512	2560	3e5	20.5	55	169	2e4	20.4	97	354	9e3
5	60	0.4	41.6	13.3%(0)	18.3	2e6	48.3	3086(4)	50.3	3e3	49.2	1728(7)	0.5%	3e3	49.2	1470(7)	1.0%	7e5	47.9	81	249	2e4	48.0	163	703	4e3
5	60	0.6	27.4	3312(1)	12.2	2e6	36.4	3341(2)	37.3	2e3	37.8	1881(6)	0.3%	3e3	37.8	1334(7)	0.5%	6e5	36.3	58	82	4e3	36.3	170	617	6e3
5	60	0.8	8.6	1.6%(0)	4.0	2e6	18.9	3572(1)	20.1	2e5	22.3	3383(2)	0.9%	5e3	22.2	2822(4)	0.9%	1e6	20.3	361	2875	2e5	20.4	787(9)	0.3%	4e4
8	40	0.4	35.4	6.4%(0)	13.8	4e6	41.7	41.5%(0)	43.1	1e4	39.4	1590	3188	1e4	39.4	679(9)	1.2%	5e5	38.5	51	202	3e4	38.5	42	165	1e4
8	40	0.6	26.5	10.1%(0)	31.2	3e6	32.2	31.9%(0)	34.1	2e4	30.3	1727(9)	0.3%	2e4	30.2	718(9)	0.4%	5e5	29.2	70	356	5e4	29.2	32	52	1e4
8	40	0.8	14.7	7.5%(0)	10.7	3e6	19.8	19.5%(0)	20.1	2e4	19.8	2750(5)	0.9%	2e4	19.8	1367(8)	0.9%	8e5	18.6	64	212	5e4	18.6	57	146	2e4
8	50	0.4	36.5	9.4%(0)	12.9	2e6	42.1	42.0%(0)	46.2	5e3	40.4	3021(3)	1.2%	1e4	40.5	3021(2)	1.9%	1e6	39.6	217	1079	1e5	39.7	179	461	5e4
8	50	0.6	27.2	11.5%(0)	25.3	2e6	32.1	32.1%(0)	33.9	5e3	31.3	0.5%(0)	1.1%	1e4	31.4	2897(4)	2.4%	2e6	30.3	277	1226	1e5	30.4	288	1233	1e5
8	50	0.8	14.9	8.8%(0)	12.6	2e6	20.1	20.0%(0)	20.4	6e3	20.1	0.4%(0)	0.9%	1e4	20.1	2353(5)	0.8%	9e5	19.0	363	1418	2e5	19.1	496	2625	2e5
8	60	0.4	36.5	12.5%(0)	14.8	1e6	41.3	41.2%(0)	49.0	2e3	39.6	0.6%(0)	1.0%	6e3	39.8	3310(1)	3.5%	1e6	38.5	296	517	9e4	38.5	358	2050	9e4
8	60	0.6	27.2	13.4%(0)	26.0	1e6	32.1	32.0%(0)	34.8	2e3	30.8	0.5%(0)	0.9%	6e3	31.0	0.9%(0)	2.3%	1e6	29.6	766(9)	0.1%	2e5	29.6	527	1470	2e5
8	60	0.8	15.7	10.4%(0)	13.7	1e6	19.8	19.8%(0)	20.7	3e3	20.1	0.5%(0)	0.7%	7e3	20.1	0.4%(0)	0.7%	1e6	18.8	1575( <b>9</b> )	0.2%	6e5	18.8	<b>1293(9</b> )	0.3%	3e5
				F <sup>G</sup>	S			$F^{kl}$	m			F <sup>ci</sup>	ıt			Fflow	/			F <sup>n</sup>	ıtz			$F^k$	m2	
Tabl	e 1.3	. Res	ults for t	he k-trimr	ned criterio	n.	_				_				_				_				_			
P	V	k	gapR	t/gap(#)	t*/gap*	nod	gapR	t/gap(#)	t*/gap*(%)	nod	gapR	t/gap(#)	t*/gap*(%)	nod	gapR	t/gap(#)	t*/gap*	nod	gapR	t/gap(#)	t*/gap*(%)	nod	gapR	t/gap(#)	t*/gap*(%)	nod
5	40	1	87.6	538(9)	1.1%	6e5	95.7	1368(8)	48.9	2e3	96.7	1803(5)	31.2	1e6	96.6	125	736	1e5	95.7	750(8)	30.3	3e5	95.7	836(8)	27.9	9e5
5	40	2	99.9	107	244	1e5	99.9	99.9%(0)	99.9	2e2	99.9	1093(7)	39.2	1e6	99.9	2358(5)	99.6%	1e6	99.9	2548(3)	95.8	6e5	99.9	2173(4)	51.6	1e6
5	50	1	96.6	<b>716(9</b> )	<b>4.9</b> %	5e5	98.8	2880(4)	89.7	2e2	99.1	1449(6)	27.2	9e5	99.1	883(8)	36.9%	3e5	98.7	1168(7)	39.3	3e5	98.7	1813(5)	32.2	7e5
5	50	2	99.9	1322( <b>8</b> )	<b>8.7</b> %	7e5	99.9	99.9%(0)	99.9	1	99.9	<b>786</b> ( <b>8</b> )	43.6	6e5	99.9	3259(1)	99.9%	6e5	99.9	2172(4)	60.3	4e5	99.9	1685(6)	62.1	8e5
5	60	1	99.9	2131(5)	0.8%	8e5	100.0	97.1%(0)	99.9	0	100.0	1482(7)	32.6	5e5	100.0	2364(5)	41.6%	8e5	99.9	<b>817</b> ( <b>8</b> )	39.5	1e5	99.9	1540(6)	34.6	5e5
5	60	2	99.9	1530( <b>8</b> )	1.3%	6e5	99.9	99.9%(0)	99.9	0	99.9	<b>1490</b> (6)	28.5	9e5	99.9	3246(1)	99.9%	9e5	99.9	1499(6)	89.9	8e4	99.9	2225(4)	70.7	4e5
8	40	2	80.5	3312(1)	3.4%	2e6	97.0	67.5%(0)	99.9	9e3	96.3	2016(5)	15.8	2e6	96.3	3451(1)	41.4%	1e6	95.9	3320(1)	35.8	2e6	95.9	<b>1846(6)</b>	6.9	2e6
8	40	3	99.9	10.0%(0)	27.5%	1e6	99.9	99.9%(0)	99.9	7e2	99.9	2618(3)	36.1	<b>4e6</b>	99.9	93.5%(0)	99.8%	1e6	99.9	91.8%(0)	99.8	1e6	99.9	<b>1901(7</b> )	21.7	1e6
8	50	2	86.3	<b>2.8</b> %(0)	5.5%	2e6	97.6	85.4%(0)	99.9	5e2	97.4	3274(2)	17.6	3e6	97.4	31.6%(0)	48.5%	1e6	97.0	38.4%(0)	59.0	1e6	97.0	2492(4)	11.8	1e6
8	50	3	99.9	47.1%(0)	67.3%	4e5	99.9	99.9%(0)	99.9	1	99.9	<b>3338</b> (1)	23.3	3e6	99.9	97.0%(0)	99.9%	4e5	99.9	95.7%(0)	100.0	4e5	99.9	3468( <b>2</b> )	19.4	1e6
8	60	2	92.1	<b>4.2</b> %( <b>0</b> )	<b>14.6</b> %	1e6	98.8	95.7%( <b>0</b> )	99.9	0	98.5	8.5%( <b>0</b> )	15.5	2e6	98.5	43.9%( <b>0</b> )	64.8%	4e5	98.1	54.8%( <b>0</b> )	65.4	6e5	98.1	7.9%( <b>0</b> )	15.8	1e6
8	60	3	99.9	85.6%(0)	98.8%	3e5	99.9	99.9%(0)	99.9	0	99.9	12.3%(0)	27.3	2e6	99.9	95.9%(0)	100.0%	2e5	99.9	96.2%(0)	99.9	2e5	99.9	<b>3201(3</b> )	22.2	8e5
				FG	<u>،</u>			F <sup>ki</sup>	m			F <sup>ci</sup>				Fflow				F <sup>n</sup>	112			F <sup>k</sup>	m2	

895

Table 2OWASTP results for  $F^{km2}$  including valid inequalities.

Table	2.1. Resu	lts for the	e k-centrum	criterion.														
P	<i>V</i>	k	gapR	<i>t/gap</i> (#)	t*/gap*	nod	gapR	<i>t/gap</i> (#)	t*/gap*	nod	gapR	<i>t/gap</i> (#)	t*/gap*	nod				
5	40	1	0.7	2.1	6.2	5e3	0.7	3.3	12	6e3	0.7	1.7	7.5	5e3				
5	40	3	0.4	1.4	6.4	2e3	0.4	1.4	5.0	2e3	0.4	0.6	1.6	1e3				
5	40	4	0.2	0.5	1.7	2e2	0.2	0.6	1.2	2e2	0.2	0.3	1.0	3e2				
5	50	1	0.6	417( <b>9</b> )	0.5%	6e5	0.6	427( <b>9</b> )	0.2%	6e5	0.6	<b>406(9</b> )	0.3%	9e5				
5	50	3	0.5	173	1636	3e5	0.5	121	1083	1e5	0.5	88	766	2e5				
5	50	4	0.2	3.2	19	2e3	0.2	2.5	12	2e3	0.2	1.0	3.9	1e3				
5	60	1	0.6	694( <b>9</b> )	<b>0.2</b> %	7e5	0.6	661( <b>9</b> )	0.4%	6e5	0.6	<b>509</b> ( <b>9</b> )	0.3%	8e5				
5	60	3	0.4	295	2536	3e5	0.4	412(9)	0.1%	4e5	0.4	323	2885	6e5				
5	60	4	0.3	8.3	57	5e3	0.3	13	101	7e3	0.3	5.0	29	5e3				
8	40	2	0.9	182	639	3e5	0.9	180	727	3e5	0.9	81	270	3e5				
8	40	4	0.7	47	256	6e4	0.7	44	276	7e4	0.7	21	106	6e4				
8	40	7	0.1	1.1	4.6	5e2	0.1	0.9	3.4	4e2	0.1	0.4	1.3	5e2				
8	50	2	0.8	1260(8)	0.9%	1e6	0.8	1487(8)	0.9%	1e6	0.8	1100(8)	0.9%	2e6				
8	50	4	0.7	839(8)	0.8%	7e5	0.7	881(8)	1.0%	6e5	0.6	<b>652</b> (9)	0.6%	1e6				
8	50	/	0.2	53	387	664	0.2	92	6//	4e4	0.2	26	177	5e4				
8	60	2	0.6	1839(7)	0.7%	166	0.6	2028(6)	0.9%	Ie6	0.6	1549(7)	0.6%	266				
8	60	4	0.5	1159(7)	0.7%	7e5	0.5	1198(7)	0.7%	7e5	0.5	1138(7)	0.6%	1e6				
8	60	/	0.1	192	1/44	le5	0.1	343 rkm2	3226	2e5	0.1	55 rkm?	4/2	6e4				
Table	2.2 Pocu	ulte for the	o Uurwicz c	ritorion	,KIII2			F ****2 +	(14)			FRANZ	+ (15)					
	2.2. Kesu		aanR	t/gan(#)	t*/aan*	nod	aanP	t /aa n(#)	t*/ga n*	nod	aanP	t/aa n(#)	t*/aa n*	nod	aanP	t/ga n(#)	t*/aa n*	nod
5	10	04	20 gupic 17 3	21	11/gup	303	47 3	10	08	203	36 6	2/gup(#)	120	203	AT 2	15	1/gup 84	203
5	40	0.4	35.0	13	40	163	35.0	10	18	002	30.0	12	27	103	35.0	72	15	2CJ 7o2
5	40	0.0	20.4	22	40	303	20.4	22	50	302	177	23	13	263	20.4	14	28	203
5	50	0.0	476	10	163	303	47.6	10	130	303	377	25	264	403	476	20	20	203
5	50	0.4	361	45	105	303	36.1	37	03	203	30.7	51	180	303	361	25	90	203
5	50	0.0	20.4	97	354	963	20.4	110	354	2c5 2e4	179	77	205	7e3	20.4	66	251	2CJ 8e3
5	60	0.0	20.4 48.0	163	703	4e3	48.0	166	409	6e3	38.9	183	700	763	48.0	148	733	5e3
5	60	0.4	36.3	170	617	6e3	36.3	100	533	7e3	31.4	156	784	6e3	363	115	733	5e3
5	60	0.0	20.4	787(9)	0.3%	4e4	20.4	850(9)	01%	1e5	18.1	825(9)	0.1%	1e5	20.4	526	2720	6e4
8	40	0.0	38.5	42	165	1e4	38.5	34	128	1e4	29.4	68	203	1e4	38.5	21	59	0c4 0e3
8	40	0.4	29.2	32	52	1e4	29.2	32	79	7e4	23.4	52	137	7e4	29.2	21	50	1e4
8	40	0.8	18.6	52	146	2e4	18.6	46	94	3e4	16.6	66	130	3e4	18.6	32	53	2e4
8	50	0.0	39.7	179	461	5e4	39.7	182	462	7e4	31.4	537	3578	5e4	39.7	122	345	4e4
8	50	0.6	30.4	288	1233	1e5	30.4	344	2168	2e5	26.2	397	1310	9e4	30.4	249	1493	1e5
8	50	0.8	191	496	2625	2e5	19.1	679(9)	0.2%	2e5	17.2	730(9)	0.2%	1e5	191	379	2262	2e5
8	60	0.4	38.5	358	2050	9e4	38.5	271	1564	9e4	30.7	599	3529	1e5	38.5	225	940	2e8 7e4
8	60	0.6	29.6	527	1470	2e5	29.7	757(9)	17.1%	3e5	25.7	1207	2819	2e5	29.6	371	712	1e5
8	60	0.8	18.8	1293( <b>9</b> )	0.3%	3e5	18.8	1537(8)	0.2%	5e5	17.1	1780(8)	0.4%	3e5	18.8	874(9)	0.2%	3e5
				F	;km2			F <sup>km2</sup> +	(14)			F <sup>km2</sup>	+ (15)			F <sup>km2</sup> +	(16)	
Table	2.3. Resu	ilts for the	e <i>k</i> -trimmed	1 criterion.														
P	V	k	gapR	t/gap(#)	t*/gap*(%)	nod	gapR	<i>t/gap</i> (#)	t*/gap*	nod	gapR	<i>t/gap</i> (#)	t*/gap*(%)	nod	gapR	t/gap(#)	t*/gap*	nod
5	40	1	95.7	836(8)	27.9	9e5	95.7	1629(6)	26.9%	1e6	79.9	1805(5)	28.2	7e5	79.7	138	1344	2e5
5	40	2	99.9	2173(4)	51.6	1e6	99.9	<b>743(8)</b>	<b>44.8</b> %	4e5	69.2	2162(4)	58.4	1e6	69.0	1088(7)	55.9%	4e5
5	50	1	98.7	1813(5)	32.2	7e5	98.7	1449(7)	26.8%	6e5	81.2	2168(4)	34.4	6e5	81.2	33	196	3e4
5	50	2	99.9	1685(6)	62.1	8e5	99.9	<b>1397</b> ( <b>7</b> )	<b>48.6</b> %	7e5	71.0	1579( <b>7</b> )	59.8	7e5	71.4	1447(6)	58.5%	3e5
5	60	1	99.9	1540(6)	34.6	5e5	99.9	2367(4)	32.5%	6e5	83.6	3248(1)	62.8	8e5	83.5	<b>813(8</b> )	8.3%	4e5
5	60	2	99.9	2225(4)	70.7	4e5	99.9	2213(4)	<b>59.5</b> %	3e5	72.7	1950( <b>5</b> )	64.2	4e5	72.6	<b>1828</b> (5)	60.8%	6e5
8	40	2	95.9	<b>1846(6)</b>	6.9	2e6	95.9	1863( <b>6</b> )	10.2%	2e6	85.0	2774(4)	15.7	2e6	85.0	1886( <b>6</b> )	7.8%	1e6
8	40	3	99.9	1901(7)	21.7	1e6	99.9	1037	3000	8e5	83.3	2384(5)	19.6	2e6	83.2	1954(5)	15.9%	1e6
8	50	2	97.0	2492(4)	11.8	1e6	97.0	<b>2170</b> (5)	11.7%	1e6	87.0	2483(4)	9.5	1e6	87.0	2256( <b>5</b> )	13.3%	7e5
8	50	3	99.9	3468( <b>2</b> )	19.4	1e6	99.9	<b>3221(2)</b>	<b>7.4</b> %	1e6	85.0	10.5%(0)	22.3	2e6	84.9	3291(1)	17.6%	2e6
8	60	2	98.1	7.9%(0)	15.8	1e6	98.1	<b>3216(2)</b>	15.9%	9e5	88.7	3575(1)	18.6	6e5	88.7	3331(1)	13.2%	8e5
8	60	3	99.9	3201( <b>3</b> )	22.2	8e5	99.9	3205( <b>3</b> )	20.8%	8e5	86.8	3304(1)	20.2	9e5	86.8	2815(3)	24.6%	6e5
				F	zkm2			$F^{km2}+$	(14)			F <sup>km2</sup>	+ (15)			$F^{km^{2}}+$	(16)	

Table 3							
OWASTP	results	for	<i>F<sup>mtz</sup></i>	including	valid	inequalities.	

Table 3.1. Results for the *k*-centrum criterion.

P	11/1	k	ganR	t/gan(#)	t*/ga n*	nod	ganR	t/ga n(#)	t*/gan*	nod	ganR	t/ga n(#)	t*/gan*	nod				
11	141	ĸ	Supr	1/gup(#)	t/gup	nou	gupk	t/gup(#)	t /gup	nou	Supr	<i>ι/gup</i> (#)	t /gup	nou				
5	40	1	0.7	3.0	7.2	3e3	0.7	4.9	13	6e3	0.7	3.1	7.3	3e3				
5	40	3	0.4	1.7	2.8	5e2	0.4	1.4	2.1	4e2	0.4	1.8	3.2	5e2				
5	40	4	0.2	0.8	2.0	8	0.2	0.9	2.1	11	0.2	0.8	2.1	8				
5	50	1	0.5	22	153	2e4	0.5	19	100	2e4	0.5	23	157	2e4				
5	50	3	0.4	4.4	16	6e3	0.4	3.4	10	3e3	0.4	4.5	16	6e3				
5	50	4	0.2	1.5	5.6	7e2	0.2	1.5	4.4	6e2	0.2	1.5	5.8	7e2				
5	60	1	0.5	75	537	8e4	0.5	270	2002	1e5	0.5	77	551	8e4				
5	60	3	0.4	39	179	3e4	0.4	26	121	5e4	0.4	40	185	3e4				
5	60	4	0.2	1.3	3.0	3e2	0.2	1.7	4.0	3e2	0.2	1.4	3.2	3e2				
8	40	2	0.9	146	372	2e5	0.9	495	1465	3e5	0.9	148	376	2e5				
8	40	4	0.7	25	126	4e4	0.7	63	152	5e4	0.7	25	128	4e4				
8	40	7	0.1	1.2	3.1	3e2	0.1	1.2	3.6	2e2	0.1	1.2	3.3	3e2				
8	50	2	0.7	849( <b>9</b> )	0.3%	6e5	0.7	1685(6)	0.4%	8e5	0.7	<b>734(9</b> )	0.2%	7e5				
8	50	4	0.5	79	503	8e4	0.5	435	2467	2e5	0.5	80	511	8e4				
8	50	7	0.1	3.3	18	2e3	0.1	2.6	5.9	6e2	0.1	3.4	19	2e3				
8	60	2	0.6	1832( <b>6</b> )	1.1%	8e5	0.6	2462(4)	1.1%	9e5	0.6	<b>1826(6)</b>	1.1%	8e5				
8	60	4	0.5	1289( <b>7</b> )	0.7%	5e5	0.5	1622( <b>7</b> )	0.5%	4e5	0.5	<b>1232(7)</b>	0.7%	6e5				
8	60	7	0.1	5.4	24	4e3	0.1	9.4	55	5e3	0.1	4.2	19	4e3				
				F <sup>mt</sup>	Z			$F^{mtz}+$	(14)			$F^{mtz}+$	(15)					
Table 3	3.2. Resu	lts for the	Hurwicz c	riterion.														
P	V	α	gapR	t/gap(#)	t*/gap*	nod	gapR	t/gap(#)	t*/gap*	nod	gapR	<i>t/gap</i> (#)	t*/gap*	nod	gapR	t/gap(#)	t*/gap*	nod
5	40	0.4	47.3	16	66	1e3	47.3	16	64	2e3	37.5	17	66	8e2	47.3	16	66	1e3
5	40	0.6	35.9	17	65	1e3	35.9	19	64	3e3	30.1	19	65	1e3	35.9	17	64	1e3
5	40	0.8	20.4	16	65	2e3	20.4	17	64	3e3	17.9	18	66	2e3	20.4	16	65	2e3
5	50	0.4	47.6	33	70	2e3	47.6	35	69	4e3	37.7	36	74	3e3	47.6	33	70	2e3
5	50	0.6	36.0	34	73	6e3	36.0	56	176	2e4	30.6	42	116	9e3	36.0	34	73	6e3
5	50	0.8	20.5	55	169	2e4	20.3	64	279	2e4	17.8	53	133	1e4	20.3	39	82	8e3
5	60	0.4	47.9	81	249	2e4	47.9	105	318	2e4	38.9	89	160	1e4	47.9	82	253	2e4
5	60	0.6	36.3	58	82	4e3	36.3	78	163	7e3	31.3	61	81	4e3	36.3	59	85	4e3
5	60	0.8	20.3	361	2875	2e5	20.3	482(9)	0.1%	1e5	18.0	345	2759	1e5	20.3	366	2919	2e5
8	40	0.4	38.5	51	202	3e4	38.5	42	121	2e4	29.4	84	300	4e4	38.5	52	204	3e4
8	40	0.6	29.2	70	356	5e4	29.2	93	424	7e4	24.5	160	969	9e4	29.2	83	477	7e4
8	40	0.8	18.6	64	212	5e4	18.6	127	723	7e4	16.6	93	297	4e4	18.6	65	214	5e4
8	50	0.4	39.6	217	1079	1e5	39.6	455	1708	1e5	32.2	445	1841	1e5	39.6	217	1093	1e5
8	50	0.6	30.3	277	1226	1e5	30.3	494	2859	2e5	26.1	483	2260	1e5	30.3	280	1244	1e5
8	50	0.8	19.0	363	1418	2e5	19.0	633	2669	2e5	17.2	700	2746	2e5	19.0	373	1499	2e5
8	60	0.4	38.5	296	517	9e4	38.5	673(9)	0.1%	1e5	30.7	1111(9)	0.1%	2e5	38.5	291	525	9e4
8	60	0.6	29.6	766(9)	0.1%	2e5	29.6	701	2928	3e5	25.6	1243(8)	0.2%	2e5	29.6	742(9)	0.1%	2e5
8	60	0.8	18.8	1575(9)	0.2%	6e5	18.8	2070(7)	0.3%	7e5	17.7	2225(6)	1.4%	5e5	18.8	1595(9)	0.2%	6e5
				F <sup>mt</sup>	z			F <sup>mtz</sup> +	(14)			F <sup>mtz</sup> +	(15)			F <sup>mtz</sup> +	(16)	
Table (	3.3. Resu	ilts for the	k-trimmed	l criterion.					()				()				()	
P	V	k	gapR	t/gap(#)	t*/gap*(%)	nod	gapR	t/gap(#)	t*/gap*(%)	nod	gapR	t/gap(#)	t*/gap*(%)	nod	gapR	t/gap(#)	t*/gap*(%)	nod
5	40	1	95.7	750(8)	30.3	3e5	95.7	1084(7)	34.9	7e5	79.8	2912(2)	55.5	1e6	79.7	1190(7)	30.1	5e5
5	40	2	99.9	2548(3)	95.8	6e5	99.9	568(9)	53.5	1e5	68.9	1096(7)	64.7	2e5	68.9	1808(5)	64.7	3e5
5	50	1	987	1168(7)	393	3e5	98.7	1467(6)	36.2	7e5	813	3332(1)	44.8	6e5	81.2	1488(6)	32.2	4e5
5	50	2	99.9	2172(4)	60.3	4e5	99.9	397(9)	19.2	5e4	71.0	1113(7)	61.5	1e5	71.0	1118(8)	476	2e5
5	60	1	99.9	817(8)	39 5	1e5	99.9	1815(5)	447	4e5	83.6	2819(3)	59.6	3e5	83.6	1546(6)	54.0	1e5
5	60	2	99.9	1499(6)	89.9	8e4	99.9	1523(6)	60.8	2e5	72.5	<b>600(9</b> )	544	4e4	72.6	1236(7)	67.8	9e4
8	40	2	95.9	3320(1)	35.8	2e6	95.9	211%(0)	361	205 3e6	85.0	13.6%(0)	24.1	2e6	85.0	3010(2)	24.0	2e6
8	40	2	99.9	91.8%(0)	99.8	1e6	99.9	2887(2)	52.8	2e6	83.3	21.9%(0)	345	1e6	83.3	25 4%(0)	35.4	9e5
8	50	2	97.0	38 4%(0)	59.0	166	970	3258(1)	40.0	166	86.0	20.3%(0)	28.5	665	870	<b>16 4</b> %(0)	<b>777</b>	8e5
8	50	2	00.0	95. <del>4</del> %(0)	100.0	100	00.0	3230(1) 31 7%(0)	54.6	106	85.0	20.3%(0) 42.0%( <b>0</b> )	20.5	105	85.0	35 5%(0)	62.3	105
8	50	נ ר	99.9 08.1	51.0%(U) 51.0%(U)	65 4	4CJ 6e5	09.9 09.1	31.7%(U) 35 5%(A)	J-1.0 /1.0	100 805	88 6 97.0	42.0%(U)	66.0	4 <del>0</del> 5	66 6 07.0	33.3%(U) 31.2%( <b>D</b> )	35.6	305
8	60	2	00.0	06 2%( <b>0</b> )	00.4	205	00.1 00.0	23.3%(U) 24.2%(A)	-11.0 573	605	90.0 86.0		74.0	205	90.0 86.0	51.2%(U) 64.5%( <b>n</b> )	78.3	205
0	00	J	33.3	30.2%( <b>U</b> )	33.3 Z	203	55.5	JH.2/0(U)	37.3 (14)	003	00.9	CO.3/0(U)	(15)	200	00.9	CH.J/6(U)	(16)	200
				F				r+	(14)			r+	(13)			r+	(10)	

 Table 4

 Comparison with the results obtained by Galand and Spanjaard (2012).

Table	e 4.1. F	Results	for the	k-centru	m criterio	n.											
P	V	k	t.	t	t*	t.	t	t*	t.	t	t*	t.	t	t*	t.	t	t*
5	20	1	0.3	11	111	1.3	11	93	0.6	11	118	0.6	0.8	1.5	0.1	0.2	0.6
5	20	3	0.2	0.8	2.0	1.4	2.3	3.5	0.2	0.8	2.1	0.0	0.5	0.9	0.0	0.1	0.2
5	20 30	4	0.1	0.4 91	2.1	0.7 14	1.0 66	1.3 229	0.0	0.3 86	2.2 334	0.0	0.1 3.1	0.6 15	0.0	0.0	0.1 4 7
5	30	3	1.2	31	593	1.0	24	225	1.0	37	833	0.7	0.8	2.0	0.1	0.4	1.1
5	30	4	0.3	12	184	4.3	5.6	9.9	0.2	4.1	64	0.0	0.3	1.1	0.0	0.1	0.3
5	40	1	-	-	-	63	628	2638	-	-	-	1.3	3.0	7.2	0.4	1.7	7.5
5	40	3	1.2	78	610	41	87	443	1.2	65	617	0.8	1.7	2.8	0.3	0.6	1.6
5	40 50	4	0.4	13	225	15	19	30	0.2	14	2/3	0.2 17	0.8 <b>22</b>	2.0 153	0.1 11	<b>0.3</b> 406	1.0 3600
5	50	3	8.5	2039	_	119	420	7360	6.1	1897	_	1.0	4.4	16	0.6	88	766
5	50	4	0.9	243	4108	46	60	103	0.5	58	366	0.2	1.5	5.6	0.1	1.0	3.9
5	60	1	-	-	-	-	-	-	-	-	-	1.8	75	537	1.5	509	3600
5	60 60	3	-	- 1116	-	62 10	660	5826	-	-	-	1.5	39	179	0.8	323	2885
5	00 70	4	0.9 -	-	-	-	-	-	5.7 -	-	-	2.5	30	5.0 92	2.2	5.0 412	29 3600
5	70	3	-	-	-	-	-	-	-	-	-	1.2	6.3	28	0.6	45	365
5	70	4	-	-	-	-	-	-	-	-	-	0.4	3.3	15	0.1	32	311
8	20	2	5.2	57	436	3.9	50	393	4.7	59	381	1.1	3.3	14	0.3	1.8	12
8	20	4	1.7	23	135	1.8	17	81	2.3	21	112	0.6	1.5	4.5	0.1	0.3	1.5
0 8	20 30	2	0.5 37	1.4	7.9 6648	25	1024	2.5	22	1269	5.1 5584	<b>0.0</b> 6.4	0.1 34	0.8 140	1.5	0.0 27	0.1 71
8	30	4	31	295	3414	15	189	1992	14	275	3109	1.2	7.1	38	0.6	5.3	20
8	30	7	0.5	3.0	26	1.5	2.6	8.4	0.1	2.3	23	0.1	0.6	1.4	0.0	0.2	0.8
8	40	2	-	-	-	-	-	-	-	-	-	6.5	146	372	7.3	81	270
8	40	4	95	1776	11820	67	1040	5029	76	1697	10283	1.0	25	126	0.5	21	106
0 8	40 50	2	-	122	-	4.4	-	-	-	78 -	-	0.2 23	1.2 849	3.1 3600	25	<b>0.4</b> 1100	3600
8	50	4	-	-	-	-	-	-	-	-	-	4.7	79	503	2.4	652	3600
8	50	7	3.5	157	1532	15	35	159	1.5	100	876	0.5	3.3	18	0.2	26	177
8	60	2	-	-	-	-	-	-	-	-	-	11	1832	3600	12	1549	3600
8	60 60	4	- 12	-	-	- วว	- 74	-	-	- 279	-	3.7	1289	3600	2.5	1138 55	3600 472
8	70	2	-	400	-	-	-	-	9.0 -	278	-	0.5 <b>41</b>	<b>J.4</b> 1919	24 3600	147	1886	472 3600
8	70	4	-	-	-	-	-	-	-	-	-	20	1268	3600	46	928	3600
8	70	7	-	-	-	33	341	3633	-	-	-	0.4	121	1043	0.1	371	3600
				$F^{GS}$			$F_{P1}^{GS}$			$F_{P2}^{GS}$			F <sup>mtz</sup>			$F^{km2} + (15)$	5)
Table	e 4.2. l	Results	for the	Hurwicz	z criterion.												
P  5	V  20	α	t.	t 12	t*	t+ 0.5	t 15	t* 26	t.	t 11	t*	t.	t 27	t*	t.	t 05	t*
5	20	0.4	0.9	1.2	2.5 3.4	1.1	2.1	2.0 3.4	0.6	1.1	2.3	0.9	2.7	9.4 5.4	0.5	0.5	0.7
5	20	0.8	0.6	1.7	4.3	0.5	1.5	3.4	0.5	1.5	3.1	1.0	3.2	7.7	0.3	0.7	1.3
5	30	0.4	2.4	4.6	10	3.6	6.0	12	2.2	4.5	10	2.9	6.0	12	1.4	2.3	3.9
5	30	0.6	1.9	8.9	42	3.1	10	44	1.6	9.1	44	2.3	4.9	8.1	0.9	5.2	23
5	30	0.8	1.5	28 19	105 51	1.2 12	19	61 57	0.5	18 19	91 45	2.3	9.4 16	34 66	1.4	6.1 15	19 84
5	40	0.4	0.J 79	46	156	12	24 52	164	77	46	<b>4</b> 5 156	7.8 8 3	10	65	4.0	7.2	04 15
5	40	0.8	6.5	71	212	7.1	53	185	4.1	52	226	7.7	16	65	9.9	14	28
5	50	0.4	22	124	324	39	143	353	22	125	336	13	33	70	10	29	70
5	50	0.6	26	367	2374	42	385	2404	26	368	2394	15	34	73	8.2	25	99
5	50 60	0.8	9.7 41	298	3665	23	218	1972	15 41	225	1979	16	55 81	169	9.7	66 149	251 722
5	60 60	0.4	41	401	4151	- 07	512	41/5	41	401	4095	20 18	58	249 82	20	140	735
5	60	0.8	6.4	1725	16779	25	867	9427	2.6	1586	29575	37	361	2875	70	526	2720
				$F^{GS}$			$F_{P1}^{GS}$			$F_{P2}^{GS}$			<i>F<sup>mtz</sup></i>			$F^{km2}$ + (10)	5)
Table	e 4.3. I	Results	for the	k-trimm	ned criterio	on.											
P	V	k	t∗	t	<i>t</i> *	t∗	t	<i>t</i> *	t.	t	t*	t.	t	t*			
5	20	1	0.5	1.4	2.4	0.5	1.5	2.4	0.8	1.1	1.9	0.5	0.6 10	<b>0.9</b>			
5 5	20 30	2	1.2 6.8	2.5 101	5.2 2187	1.2 6.8	∠.⊃ 102	5.4 2209	0./ 12	1.0 1441	1.9 3600	<b>U.4</b> 13	10	3.0			
5	30	2	8.6	468	9778	8.7	434	8720	1.3	42	<b>392</b>	0.6	1081	3600			
5	40	1	9.5	1160	12874	9.6	1120	13655	1.9	1084	3600	3.2	138	1344			
5	40	2	20	168	1029	20	169	1039	2.1	568	3600	2.4	1088	3600			
5	50	1	29	733	15269	29	732	15239	5.8	1467	3600	4.7	33	<b>196</b>			
5 5	50 60	2	49 45	191 370	1215	49 46	191 370	1214	4.0 62	397 1815	3600	1.5 0.1	144/ 812	3600			
5	60	2	83	829	6407	83	828	6383	19	1523	3600	5.5	1828	3600			
	-		-	$F^{GS}$	-	-	$F_{P1}^{GS}$		-	$F^{mtz}$ + (1	4)		$F^{km2} + (1)$	6)			

Table 5				
OWASTP	results	for	large	instances.

Table 5.	1. Results for	the k-centrum	criterion.							
P	V	k	gapR	<i>t/gap</i> (#)	t*/gap*	nod	gapR	<i>t/gap</i> (#)	t*/gap*	nod
5	40	1	0.7	3.0	7.2	3e3	0.7	1.7	7.5	5e3
5	40	3	0.4	1.7	2.8	5e2	0.4	0.6	1.6	1e3
5	40	4	0.2	0.8	2.0	8	0.2	0.3	1.0	3e2
э 5	50	1	0.5	22	153	2e4 6e3	0.6	406(9)	0.3%	9e5 2e5
5	50	4	0.4	15	56	0e3 7e2	0.3	10	39	263 1e3
5	60	1	0.5	75	537	8e4	0.6	509(9)	0.3%	8e5
5	60	3	0.4	39	179	3e4	0.4	323	2885	6e5
5	60	4	0.2	1.3	3.0	3e2	0.3	5.0	29	5e3
5	80	1	0.3	230	1033	8e4	0.4	1453(6)	0.3%	1e6
5	80	3	0.2	18	148	7e3	0.3	846(8)	0.2%	7e5
5	80	4	0.1	2.0	6.4 0.2%	1e2	0.1	19	95	1e4
5 5	100	1	0.2	750(8) 188	0.2%	30/	0.3	568(9)	1.0%	365
5	100	4	0.1	6.6	45	7e2	0.2	130	1186	4e4
8	40	2	0.9	146	372	2e5	0.9	81	270	3e5
8	40	4	0.7	25	126	4e4	0.7	21	106	6e4
8	40	7	0.1	1.2	3.1	3e2	0.1	0.4	1.3	5e2
8	50	2	0.7	<b>849(9</b> )	0.3%	6e5	0.8	1100(8)	0.9%	2e6
8	50	4	0.5	79	503	8e4	0.6	652(9)	0.6%	1e6
8	50	/	0.1	3.3 1922(6)	18	2e3	0.2	26 1540(7)	1//	5e4 2o6
8	60	4	0.5	1289( <b>7</b> )	0.7%	5e5	0.5	1138(7)	0.6%	2e0 1e6
8	60	7	0.1	5.4	24	4e3	0.1	55	472	6e4
8	80	2	0.3	2520(5)	<b>0.2</b> %	2e6	0.4	3281(1)	0.8%	2e6
8	80	4	0.3	293	1449	2e5	0.3	1292(8)	0.6%	1e6
8	80	7	0.1	3.5	13	9e2	0.1	379(9)	0.3%	2e5
8	100	2	0.4	<b>3591</b> (1)	0.4%	9e5	0.4	0.3%(0)	1.1%	1e6
8	100	4	0.3	2606(4)	0.3%	7e5	0.3	2944(3)	0.8%	166
o 10	40	2	0.1	17 2229(5)	82 0.4%	165 2e6	10	1062(9)	0.2%	904 2e6
10	40	4	0.8	470	949	6e5	0.8	267	1872	2e0 7e5
10	40	7	0.4	7.9	27	9e3	0.4	14	109	4e4
10	50	2	0.8	3336(1)	0.5%	2e6	0.9	<b>2910(4</b> )	0.7%	4e6
10	50	4	0.7	2651(5)	0.3%	1e6	0.7	<b>1323(8</b> )	0.6%	2e6
10	50	7	0.4	142	481	7e4	0.5	746(9)	0.3%	1e6
10	60	2	0.7	<b>0.4</b> %( <b>0</b> )	0.7%	1e6	0.7	0.4%(0)	0.9%	4e6
10 10	60	4	0.6	3331(2) 872(0)	0.5%	266	0.6	2273(5) 1499(7)	0.6%	266
10	80	2	0.3	0.5%(0)	11%	4e5	0.4	<b>04%(0)</b>	0.3%	2e0 2e6
10	80	4	0.5	0.3%(0)	0.7%	9e5	0.4	3289(2)	0.5%	2e6
10	80	7	0.3	2403(5)	0.5%	4e5	0.3	<b>2014(6)</b>	0.5%	1e6
10	100	2	0.5	<b>0.4</b> %(0)	0.7%	5e5	0.5	<b>3519</b> (1)	1.2%	1e6
10	100	4	0.4	3265(1)	0.6%	5e5	0.4	<b>3259</b> (1)	1.2%	9e5
10	100	7	0.2	<b>2119</b> (5)	<b>0.3</b> %	2e5	0.3	2358(4)	0.6%	7e5
Table 5	2 Results for	the Hurwicz (	riterion	1	-m.z			F	+ (15)	
	2. ICSUITS 101		ganR	t/gan(#)	t*/ga n*	nod	ganR	t/gan(#)	t*/ga n*	nod
5	40	0.4	47.3	16	66	1e3	47.3	15	84	2e3
5	40	0.6	35.9	17	65	1e3	35.9	7.2	15	7e2
5	40	0.8	20.4	16	65	2e3	20.4	14	28	2e3
5	50	0.4	47.6	33	70	2e3	47.6	29	<b>70</b>	2e3
5	50	0.6	36.0	34	73	6e3	36.1	25	99 251	2e3
5 5	5U 60	0.8	20.5 <b>47 0</b>	55 81	109 240	204 2e4	<b>20.4</b> 48.0	טט 148	201 733	863 563
5	60	0.4	36.3	58	2-13 82	4e3	36.3	115	748	5e3
5	60	0.8	20.3	361	2875	2e5	20.4	526	2720	6e4
5	80	0.4	47.4	295	1201	5e4	47.7	711(9)	0.4%	9e3
5	80	0.6	32.5	148	307	1e4	32.5	480	2446	1e4
5	80	0.8	20.2	259	1164	3e4	20.2	1066(8)	0.2%	3e4
5	100	0.4	47.8	825(9)	0.2%	1e5	47.8	945( <b>9</b> )	<b>0.2</b> %	8e3
5	100	0.6	36.2	196	532 0.0°	1e4	36.2	10/6(8)	0.1%	6e3
5	100 40	0.8	20.1	<b>ສວ4(ð</b> ) 51	0.0% 202	105 3e4	20.1	1/13(/) <b>21</b>	U.3% 59	∠€4 9e3
8	40	0.4	29.2	70	356	5e4	29.2	23	50	1e4
8	40	0.8	18.6	64	212	5e4	18.6	32	53	2e4
8	50	0.4	39.6	217	1079	1e5	39.7	122	345	4e4
8	50	0.6	30.3	277	1226	1e5	30.4	249	1493	1e5
8	50	0.8	19.0	363	1418	2e5	19.1	379	2262	2e5
8	60	0.4	38.5	296	517 0.1%	9e4	38.5	225	940	7e4
8 8	60 60	0.6	29.6 18 8	700(9) 1575( <b>9</b> )	0.1%	205 6e5	29.6 18 s	371 874(9)	0.2%	100 365
0	00	0.0	10.0	1313(3)	0.2/0	003	10.0	01-1(3)	0.4/0	303

(continued on next page)

Table 5	(continued)
---------	-------------

P	V	k	gapR	<i>t/gap</i> (#)	t*/gap*	nod	gapR	<i>t/gap</i> (#)	t*/gap*	nod
8	80	0.4	38.5	1823(7)	0.4%	3e5	38.6	<b>1375</b> (8)	0.7%	2e5
8	80	0.6	29.6	1761(7)	0.2%	3e5	29.6	<b>1514(8)</b>	0.1%	4e5
8	80	0.8	18.8	1611	3032	3e5	18.8	1119	2357	3e5
8	100	0.4	38.7	2758(4)	0.5%	2e5	38.1	<b>2313(6)</b>	0.4%	2e5
8	100	0.6	29.7	2733( <b>4</b> )	0.5%	2e5	29.7	<b>2653</b> (4)	0.2%	3e5
8	100	0.8	18.9	3449(1)	0.3%	3e5	19.0	<b>3005(4</b> )	0.7%	4e5
10	40	0.4	35.8	154	786	1e5	35.8	82	352	6e4
10	40	0.6	27.2	320	1028	2e5	27.2	150	443	1e5
10	40	0.8	17.5	545	1936	5e5	17.5	435	1158	4e5
10	50	0.4	36.0	1012(8)	0.2%	5e5	36.0	454	2961	2e5
10	50	0.6	27.5	1541(7)	0.2%	7e5	28.9	705	2695	4e5
10	50	0.8	18.0	2618(5)	0.4%	1e6	18.0	<b>2191(6)</b>	0.3%	1e6
10	60	0.4	35.7	1847(7)	0.2%	6e5	35.7	<b>1361(8)</b>	0.2%	6e5
10	60	0.6	27.1	2622(5)	0.5%	8e5	27.2	<b>2247(9)</b>	0.4%	8e5
10	60	0.8	17.7	3544(1)	0.5%	1e6	17.7	<b>3439(2)</b>	0.3%	2e6
10	80	0.4	35.0	3388(2)	0.9%	4e5	35.0	<b>2743</b> ( <b>4</b> )	0.4%	6e5
10	80	0.6	27.0	3448(1)	0.3%	4e5	27.0	3123(3)	0.4%	6e5
10	80	0.8	17.7	0.4%(0)	0.8%	4e5	17.7	0.2%(0)	0.6%	8e5
10	100	0.4	35.0	0.3%(0)	0.7%	2e5	34.9	0.2%(0)	0.4%	4e5
10	100	0.6	26.9	0.3%(0)	0.8%	2e5	26.8	0.2%(0)	0.4%	4e5
10	100	0.8	17.6	0.3%(0)	0.5%	2e5	17.6	0.2%(0)	0.4%	4e5
					Fmtz			F <sup>km2</sup>	$^{2}+(16)$	



Fig. 5. TG performance profiles for each instance size (|P|, |V|), k-centrum criterion and some selected formulations:  $F^{CS}$  (dotted line),  $F^{ntz}$  (solid line),  $F^{km2}$  + (15) (dashed line).

tively. Note that this index has been scaled on the *y*-axis in order to illustrate a measure with the time and gap obtained. We focus on the comparison with the formulation that already exists in the literature,  $F^{GS}$ , and those with a better performance in our study. From Fig. 5(a) we conclude that the best performance for OWASTP is obtained with formulations  $F^{mtz}$  and  $F^{km2} + (15)$ , being  $F^{km2} + (15)$  slightly outperformed by  $F^{mtz}$ . In terms of TG index, Fig. 5(b) shows that  $F^{mtz}$  is more stable with 5 objectives but the maximum TG value is stabilized for instances of 8 objectives and  $|V| = \{50, 60\}$ .

Table 4.1 shows that, in nearly all cases,  $F_{P1}^{mtz}$  and  $F_{P2}^{km2}$ + (16) have a better performance than  $F^{GS}$ ,  $F_{P1}^{GS}$  and  $F_{P2}^{GS}$  (Galand & Spanjaard, 2012).

Table 5.1 shows our results with formulations  $F^{mtz}$  and  $F^{km2}$ + (16) for larger graphs of sizes up to 100 nodes and with up to 10 objectives. These instances are larger than the largest ones reported so far in the literature with the *k*-centrum criterion, which, to the best of our knowledge have up to 60 nodes and up to 8 objectives (Galand & Spanjaard, 2012). We can observe that, when  $|V| \ge 80$ , after 1 hour of computing time there are already some unsolved instances. Nevertheless, the performance of  $F^{mtz}$  and  $F^{km2}$ + (16) is remarkable, as the biggest gaps at termination are always around 1%.

## 4.2. Tests with the Hurwicz criterion

In Table 1.2, the results of block  $F^{GS}$  indicate that the OWASTP formulation of Galand and Spanjaard (2012) produces the smallest gaps at the root node (gapR), although only few instances could be solved to optimality, and the gaps remaining at termination (gap\*) are outperformed by most of the other formulations. The results of block *F*<sup>km</sup> show that the number of instances solved to optimality is higher than that of  $F^{GS}$ , although the gaps in the unsolved instances are higher. We recall that  $F^{km}$  uses  $O(n^3)$  variables and constraints, which can be too high in large graphs. This also explains the low average number of explored nodes in the B&B tree. The results of block F<sup>cut</sup> indicate that, in general, this formulation outperforms both  $F^{GS}$  and  $F^{km}$ . As can be seen in block  $F^{flow}$  this formulation improves the average running times and gaps of F<sup>cut</sup> but the maximum optimality gaps at termination are still competitive in  $F^{cut}$  against  $F^{flow}$ . The blocks  $F^{mtz}$  and  $F^{km2}$  show the best performance in terms of number of instances solved to optimality, running times and optimality gaps.

Table 2.2 shows that the performance of  $F^{km2}$  is improved when constraints (16) are added. On the contrary, Table 3.2 shows that the performance of  $F^{mtz}$  is almost not improved adding any of these valid inequalities.



Fig. 6. TG performance profiles for each instance size (|P|, |V|), Hurwicz criterion and some selected formulations: F<sup>GS</sup> (dotted line), F<sup>mtz</sup> (solid line), F<sup>km2</sup> + (16) (dashed line).

Fig. 6 summarizes, in terms of the TG index, the main results of Tables 1.2, 2.2 and 3.2. From Fig. 6(a) we conclude that the best performance for OWASTP is obtained with formulations  $F^{mtz}$  and  $F^{km2}$ + (16), being  $F^{mtz}$  slightly outperformed by  $F^{km2}$ + (16). In terms of maximum TG index, Fig. 6(b) shows the best performance for  $F^{mtz}$  and  $F^{km2}$ + (16).

In Table 4.2 our results in columns  $F^{mtz}$  and  $F^{km2}$ + (16) show a better performance as compared with the results reported in Galand and Spanjaard (2012) in almost all cases. Table 4.2 shows that, in nearly all cases,  $F^{mtz}$  and  $F^{km2}$ + (16) have a better performance than  $F^{CS}$ ,  $F_{P1}^{CS}$  and  $F_{P2}^{CS}$ .

Table 5.2 shows our results with formulations  $F^{mtz}$  and  $F^{km2}$ + (16) with respect to the Hurwicz criterion with larger graphs of sizes up to 100 nodes and with up to 10 objectives. To the best of our knowledge the largest OWASTP instances reported so far in the literature with this criterion have up to 60 nodes and up to 5 objectives. We can observe that, when  $|V| \ge 80$ , after 1 hour of computing time there are still some unsolved instances. Nevertheless, the performance of  $F^{mtz}$  and  $F^{km2}$ + (16) is remarkable, as the biggest gaps at termination are always below 1%.

## 4.3. Tests with the k-trimmed criterion

The results of the *k*-trimmed criterion indicate that this is the hardest objective function as compared to the *k*-centrum and Hurwicz. This difficulty lies on the fact that the LP bound at the root node of the B&B tree is rather poor (zero in nearly all cases) which produces percentage optimality gaps at the root node (*gapR*) of almost 100% for all formulations without reinforcements.

Table 1.3 shows that for |P| = 5 the best results in terms of average and maximum gaps at termination are obtained with  $F^{GS}$ . In contrast the results of block  $F^{km}$  indicate that solving the LP relaxation of the instances in less than 1 hour of computing time, becomes nearly impossible (observe that, in general, the number of explored nodes in the B&B tree is very small). Once again, we recall that  $F^{km}$  uses  $O(n^3)$  variables and constraints, which can be too demanding for large graphs. The results of block  $F^{cut}$  indicate that, in general, this formulation outperforms both  $F^{GS}$  and  $F^{km}$  for |P| = 8. As can be seen, formulations  $F^{flow}$  and  $F^{mtz}$  exhibit a similar performance, which is not competitive against  $F^{GS}$  and  $F^{km2}$ . Formulation  $F^{km2}$  shows the best performance for |P| = 8 in terms of number of instances solved to optimality, running times and optimality gaps, and the second best for |P| = 5, just after  $F^{GS}$ .

Table 2.3 shows that the performance of  $F^{km2}$  for |P| = 8 improves when constraints (14) are added to strengthen the formulation. On the contrary, the best performance for  $F^{km2}$  and |P| = 5 is obtained for  $F^{km2} + (16)$ . We can observe that both  $F^{GS}$  and  $F^{km2} + (16)$  produce the best results for the *k*-trimmed criterion and |P| = 5. Table 3.3 shows that the performance of  $F^{mtz}$  only im-

proves when constraints (14) are added, but this improvement is not enough to outperform  $F^{GS}$  or  $F^{km2}$  + (16).

Fig. 7 summarizes in terms of the TG index, the main results of Tables 1.3, 2.3 and 3.3 as follows. From Fig. 7(a) we conclude that the best performance for OWASTP and |P| = 5 is obtained with formulations  $F^{GS}$  and  $F^{km2}$ + (16), being  $F^{km2}$ + (16) slightly outperformed by  $F^{GS}$ . When the number of objectives increases to |P| = 8, the best performance is attained by  $F^{km2}$ + (16). Similarly, with respect to worse cases (Fig. 7(b)),  $F^{GS}$  produces the best the TG index for |P| = 5 and  $F^{km2}$ + (16) for |P| = 8.

Finally, we observe in Table 4.3 that, in contrast with the results obtained for the *k*-centrum and Hurwicz criteria, formulations  $F^{mtz}$ + (14) and  $F^{km2}$ + (16) do not outperform the results reported by Galand and Spanjaard (2012) for the *k*-trimmed criterion. The reader may note that already for |P| = 8 there are several *k*-trimmed instances which could not be solved in one hour (see Table 4.3). This shows that |P| = 8 is the size limit of the instances that can be solved in one hour. This has been the reason for not including a third table with larger instances for the *k*-trimmed criterion.

### 4.4. Performance summary

In summary, we observe that the performance of the OWAST formulations largely depend on the choice for the objective function (that is, the weights vector). In particular we conclude, from our computational experience, that for the *k*-centrum and  $|P| \in \{5, ..., N\}$ 8}, it is convenient to apply formulations  $F^{mtz}$  and  $F^{km2}$ + (15), being  $F^{km2}$ + (15) slightly outperformed by  $F^{mtz}$ . When |P| = 10 the performance of both  $F^{mtz}$  and  $F^{km2}$  + (16) is remarkable, as the biggest gaps at termination are always around 1%. The conclusion for the Hurwicz criterion is different, since the best performance for OWASTP is obtained with formulations  $F^{mtz}$  and  $F^{km2}$ + (16), being  $F^{mtz}$  slightly outperformed by  $F^{km2}$ + (16). When |P| = 10the performance of both  $F^{mtz}$  and  $F^{km2}$ + (16) is remarkable, as the biggest gaps at termination are always around 1%. However, in this case,  $F^{mtz}$  is outperformed by  $F^{km2}$  + (16). Finally, we conclude that the best performance for the k-trimmed criterion and |P| = 5 is obtained with formulations  $F^{GS}$  and  $F^{km2}$  + (16), being  $F^{km2}$ + (16) slightly outperformed by  $F^{GS}$ . When the number of objectives increases to |P| = 8, the best performance is attained by  $F^{km2}$ + (16). Therefore, we cannot conclude that a specific formulation is superior to all the others regardless of the objective function considered. This reinforces the interest of the catalog of formulations and valid inequalities, and the extensive numerical results from computational experiments presented in this paper.



Fig. 7. TG performance profiles for each instance size (|P|, |V|), k-trimmed criterion and some selected formulations: FGS (line dotted), Fmtz (line solid), Fmtz (line dashed).

## 5. Conclusions

In this paper we have presented reinforced mathematical programing formulations for OWASTP as well as alternative new formulations which reduces the number of decision variables. These new formulations reinforced with appropriate constraints have shown to be very effective for efficiently solving medium size OWASTP instances. However, from the obtained results it is also clear that for solving larger OWASTP instances with more objective functions further improvements are needed. Our current research focuses on the design of more sophisticated elimination tests as well as on alternative formulations leading to tighter LP bounds.

#### Acknowledgments

The research of the first author has been partially supported by the Spanish Ministry of Economy and Competitiveness through grants MECD-JCASTILLEJO PRX15/00086 and MTM2015-63779-R (MINECO/FEDER). The second, third and fourth authors were partially supported by the projects FQM-5849 (Junta de Andalucía\FEDER) and MTM2013-46962-C02-01 and MTM2016-74983-C02-01 (MINECO/FEDER). This support is gratefully acknowledged.

## References

- Andersen, K., Jörnsten, K., & Lind, M. (1996). On bicriterion minimal spanning trees: an approximation. Computers & Operations Research, 23(12), 1171-1182.
- Averbakh, I. (2001). On the complexity of a class of combinatorial optimization problems with uncertainty. Mathematical Programming, 90, 263-272.
- Averbakh, I., & Berman, O. (1995). Probabilistic sales-delivery man and sales-delivery facility location problems on a tree. Transportation Science, 29, 184-197.
- Balas, E., & Padberg, M. (1972). On the set-covering problem. Operations Research, 20, 1152-1161.
- Boland, N., Domínguez-Marín, P., Nickel, S., & Puerto, J. (2006). Exact procedures for solving the discrete ordered median problem. Computers & Operations Research, 33(11), 3270-3300.
- Brandstadt, A., Le, V., & Spinrad, J. (2000). Graph classes: A survey. Philadelphia, PA: Society for Industrial and Applied Mathematics (SIAM)
- Breuer, M. A. (1970). Simplification of the covering problem with application to Boolean expressions. Journal of the Association for Computing Machinery, 17, 166-181.
- Calvete, H., & Mateo, P. (1998). Lexicographic optimization in generalized network flow problems. Journal of Operations Research Society, 49(2), 519-529.
- Christofides, N., & Korman, S. (1974). A computational survey of methods for the set covering problem. Management Science, 21(5), 591-599.
- Conde, E. (2004). An improved algorithm for selecting p items with uncertain returns according to the minmax-regret criterion. Mathematical Programming, 100, 345-353.
- Croce, F. D., Paschos, V. T., & Tsoukias, A. (1999). An improved general procedure for lexicographic bottleneck problems. Operations Research Letters, 24, 187–194.
- Edmonds, J. (1970). Submodular functions, matroids, and certain polyhedra. In Combinatorial structures and their applications (pp. 69-87). New York: Gordon and Breach.
- Edmonds, J. (1971). Matroids and the greedy algorithm. Mathematical Programming, 1. 127-136.
- Ehrgott, M. (2005). Multicriteria optimization. Berlin Heidelberg: Springer.

- Espejo, I., Marin, A., Puerto, J., & Rodríguez-Chía, A. (2009). A comparison of formulations and solution methods for the minimum-envy location problem. Computers & Operations Research, 36, 1966-1981.
- Fernández, E., Pozo, M. A., & Puerto, J. (2013). A modeling framework for ordered weighted average combinatorial optimization. arXiv:1306.1426v1.
- Fernández, E., Pozo, M. A., & Puerto, J. (2014). Ordered weighted average combinatorial optimization: Formulations and their properties. Discrete Applied Mathematics, 169(0), 97–118.
- Galand, L., Perny, P., & Spanjaard, O. (2010). Choquet-based optimisation in multiobjective shortest path and spanning tree problems. European Journal of Operational Research, 204(2), 303-315.
- Galand, L., & Spanjaard, O. (2012). Exact algorithms for OWA-optimization in multiobjective spanning tree problems. Computers & Operations Research, 39, 1540-1554.
- Galil, Z., & Schieber, B. (1998). On finding most uniform spanning trees. Discrete Applied Mathematics, 20, 173–175.
- Garey, M., & Johnson, D. (1979). Computers and intractability: A guide to the theory of NP-completeness. New York, NY, USA: W.H. Freeman and Co.
- Garfinkel, R., Fernández, E., & Lowe, T. (2006). The k-centrum shortest path problem. TOP. 14. 279-292.
- Gavish, B. (1983). Formulations and algorithms for the capacitated minimal directed tree problem. Journal of the ACM, 30, 118-132.
- Gouveia, L. (1995). Using the Miller-Tucker-Zemlin constraints to formulate a minimal spanning tree problem with hop constraints. Computers & Operations Research. 22(9), 959-970.
- Grzybowski, J., Nickel, S., Pallaschke, D., & Urbanski, R. (2011). Ordered median functions and symmetries. Optimization, 60(7), 801-811.
- Gupta, M., Gupta, Y., & Bector, C. (1990). Minimizing the flow-time variance in single-machine systems. Journal of Operations Research Society, 41(8), 769-779.
- Gupta, S. K., & Punnen, A. P. (1988). Minimum deviation problems. Operations Research Letters, 7(4), 201-204.
- Gusfield, D. (1990). Very simple methods for all pairs network flow analysis. SIAM
- Journal on Computing, 19(1), 143–155. Hamacher, H., & Ruhe, G. (1994). On spanning tree problems with multiple objectives. Annals of Operations Research, 52, 209-230.
- Hansen, P. (1979). Bicriterion path problems. In Multiple criteria decision making theory and application. In Lecture notes in economics and mathematics systems: Vol. 177 (pp. 109-127). Berlin: Springer.
- Hansen, P., & Labbé, M. (1988). Algorithms for voting and competitive location on a network. Transportation Science, 22(4), 278-288.
- Hansen, P., Labbé, M., & Thisse, J. (1991). From the median to the generalized center. RAIRO Operations Research, 25, 73-86.
- Hu, T. (1974). Optimum communication spanning trees. SIAM Journal on Computing, 3. 188-195.
- Hurwicz, L. (1951). Optimality criteria for decision making under ignorance p. 370. Chicago: Cowles Commission Discussion Paper
- Kalcsics, J., Nickel, S., & Puerto, J. (2003). Multi-facility ordered median problems on networks: A further analysis. Networks, 41, 1-12.
- Kalcsics, J., Nickel, S., Puerto, J., & Tamir, A. (2002). Algorithmic results for ordered median problems defined on networks and the plane. Operations Research Letters, 30, 149-158.
- Karp, R. (1972). Reducibility among combinatorial problems (pp. 85-103). New York: Plenum Press.
- Kasperski, A., & Zielinski, P. (2015). Combinatorial optimization problems with uncertain costs and the OWA criterion. Theoretical Computer Science, 565, 102-112.
- Kelly, L. M. (1944). Covering problems. National Mathematics Magazine, 19, 123-130. Kruskal, J. (1956). On the shortest spanning subtree of a graph and the traveling
- salesman problem. Proceedings of the American Mathematical Society, 7, 48-50. Landete, M., & Marín, A. (2014). Looking for edge-equitable spanning trees. Comput-
- ers & Operations Research, 41(0), 44-52. Laporte, G. (1992). The traveling salesman problem: An overview of exact and ap-
- proximate algorithms. European Journal of Operational Research, 59(2), 231-247.
- Lawler, E. L. (1966). Covering problems: Duality relations and a new method of solution. SIAM Journal on Applied Mathematics, 14, 1115-1132.

- Lawler, E. L. (1972). A procedure for computing the k best solutions to discrete optimization problems and its application to the shortest path problem. *Management Science*, 18(7), 401–405.
- Magnanti, T. L., & Wolsey, L. A. (1995). Optimal trees. In C. Ball, T. Magnanti, & G. Nemhauser (Eds.), Network models: Handbooks in operations research and management science: Vol. 7 (np. 503–615). Elsevier.
- management science: Vol. 7 (pp. 503–615). Elsevier.
  Marín, A., Nickel, S., Puerto, J., & Velten, S. (2009). A flexible model and efficient solution strategies for discrete location problems. *Discrete Applied Mathematics*, 157(5), 1128–1145.
- Martello, S., Pulleyblank, W., Toth, P., & de Werra, D. (1984). Balanced optimization problems. *Operations Research Letters*, 3(5), 275–278.
- Martin, R. (1991). Using separation algorithms to generate mixed integer model reformulations. Operations Research Letters, 10(3), 119–128.
- Mesa, J., Puerto, J., & Tamir, A. (2003). Improved algorithm for several network location problems with equality measures. *Discrete Applied Mathematics*, 130, 437–448.
- Miller, C. E., Tucker, A. W., & Zemlin, R. A. (1960). Integer programming formulation of traveling salesman problems. *Journal of the ACM*, 7(4), 326–329.
- Minoux, M. (1989). Network synthesis and optimum network design problems: models, solution methods and applications. *Networks*, 19, 313–360.
- López de los Mozos, M., Mesa, J., & Puerto, J. (2008). A generalized model of equality measures in network. *Computers & Operations Research*, 35, 651–660.
- Ogryczak, W., & Olender, P. (2012). On MILP models for the OWA optimization. Journal of Telecommunications and Information Technology, 2(2012), 5–12.
- Ogryczak, W., & Sliwinski, T. (2003). On solving linear programs with the ordered weighted averaging objective. European Journal of Operational Research, 148(1), 80–91.
- Ogryczak, W., & Tamir, A. (2003). Minimizing the sum of the k largest functions in linear time. *Information Processing Letters*, 85(3), 117–122.
- Pascoal, M., Captivo, M. E., & Clímaco, J. (2003). A note on a new variant of Murty's ranking assignments algorithm. 40R, 1(3), 243–255.
- Prim, R. (1957). Shortest connection networks and some generalizations. Bell System Technical Journal, 36, 1389–1401.
- Puerto, J. (2008). A new formulation of the capacitated discrete median problem with {0, 1}-assignment. In J. Kalcsics, & S. Nickel (Eds.), *Proceedings of the 2007* on operations research (pp. 165–170). Heidelberg: Springer.
- Puerto, J., & Rodríguez-Chía, A. (2003). Robust positioning of service units. Stochastic Models, 19(1), 125–147.

- Puerto, J., & Rodríguez-Chía, A. (2015). Ordered median location problems. In G. Laporte, S. Nickel, & F. S. d. Gama (Eds.), *Location science* (pp. 249–288). Heidelberg: Springer.
- Puerto, J., & Tamir, A. (2005). Locating tree-shaped facilities using the ordered median objective. *Mathematical Programming*, 102(2), 313–338.
- Punnen, A. (1992). Traveling salesman problem under categorization. Operations Research Letters, 12(2), 89–95.
- Punnen, A., Nair, K., & Aneja, Y. (1995). Generalized bottleneck problems. Optimization, 35(2), 159–169.
- Punnen, A. P. & Aneja, Y. P. (1997). Minimum dispersion problems. Discrete Applied Mathematics, 75(1), 93–102.
- Ramos, R., Alonso, S., Sicilia, J., & González, C. (1998). The problem of the optimal biobjective spanning tree. European Journal of Operational Research, 111(3), 617–628.
- Richey, M. (1990). Optimal location of a path or tree on a network with with cycles. *Networks*, 20, 391–407.
- Richey, M., & Punnen, A. (1992). Minimum perfect bipartite matchings and spanning trees under categorization. *Discrete Applied Mathematics*, 39, 147–153.
- Schrijver, A. (1983). Min-max results in combinatorial optimization. In mathematical programming: The state of the art (Bonn, 1982) (pp. 439–500). Berlin: Springer.
- Slater, P. (1978a). Centers to centroids in graphs. Journal of Graph Theory, 2(3), 209–222.
- Slater, P. (1978b). Structure of the k-centra in a tree. In Proceedings of the ninth Southeastern conference on combinatorics, graph theory, and computing: Vol. XXI (pp. 663–670).
- Sourd, F., & Spanjaard, O. (2008). A multi-objective branch-and-bound framework. application to the bi-objective spanning tree problem. *INFORMS Journal of Computing*, 20(3), 472–484.
- Steiner, S., & Radzik, T. (2008). Computing all efficient solutions of the biobjective minimum spanning tree problem. Computers & Operations Research, 35(1), 198–211.
- Tamir, A. (2000). The k-centrum multi-facility location problem. Dis, 109, 292–307.
- Tamir, A., Puerto, J., & Perez-Brito, D. (2002). The centdian subtree on tree networks. Discrete Applied Mathematics, 118, 263–278.
- Yen, J. Y. (1971). Finding the K shortest loopless paths in a network. Management Science, 17, 712–716.
- Yu, G. (1998). Min-max optimization of several classical discrete optimization problems. Journal of Optimization Theory and Applications, 98(1), 221–242.