

Full length article

Time and dose dependent effects of oxidative stress induced by cumene hydroperoxide in neuronal excitability of rat motor cortex neurons



R. Pardillo-Díaz^a, L. Carrascal^a, M.F. Muñoz^b, A. Ayala^b, P. Nunez-Abades^{a,*}

^a Department of Physiology, School of Pharmacy, University of Seville, Spain

^b Department of Biochemistry and Molecular Biology, School of Pharmacy, University of Seville, Spain

ARTICLE INFO

Article history:

Received 30 June 2015

Received in revised form 22 January 2016

Accepted 9 February 2016

Available online 11 February 2016

Keywords:

Amyotrophic Lateral Sclerosis

Cumene hydroperoxide

Lipid peroxidation

Neuronal excitability

Patch clamp

Pyramidal neurons

ABSTRACT

It has been claimed that oxidative stress and the production of reactive oxygen radicals can contribute to neuron degeneration and might be one factor in the development of different neurological diseases. In our study, we have attempted to clarify how oxidative damage induces dose dependent changes in functional membrane properties of neurons by means of whole cell patch clamp techniques in brain slices from young adult rats. Our research demonstrates physiological changes in membrane properties of pyramidal motor cortex neurons exposed to 3 concentrations of cumene hydroperoxide (CH; 1, 10 and 100 μM) during 30 min. Results show that oxidative stress induced by CH evokes important changes, in a concentration and time dependent manner, in the neuronal excitability of motor cortex neurons of the rat: (i) Low concentration of the drug (1 μM) already blocks inward rectifications (sag) and decreases action potential amplitude and gain, a drug concentration which has no effects on other neuronal populations, (ii) 10 μM of CH depresses the excitability of pyramidal motor cortex neurons by decreasing input resistance, amplitude of the action potential, and gain and maximum frequency of the repetitive firing discharge, and (iii) 100 μM completely blocks the capability to produce repetitive discharge of action potentials in all cells. Both larger drug concentrations and/or longer times of exposure to CH narrow the current working range. This happens because of the increase in the rheobase, and the reduction of the cancellation current. The effects caused by oxidative stress, including those produced by the level of lipid peroxidation, are practically irreversible and, this, therefore, indicates that neuroprotective agents should be administered at the first symptoms of alterations to membrane properties. In fact, the pre-treatment with melatonin, acting as an antioxidant, prevented the lipid peroxidation and the physiological changes induced by CH. Larger cells (as estimated by their cell capacitance) were also more susceptible to oxidative stress. Our results provide previously unavailable observations that large size and high sensitivity to oxidative stress (even at low concentrations) make pyramidal neurons of the motor cortex, in particular corticofugal neurons, more susceptible to cell death when compared with other neuronal populations. These results could also shed some light on explaining the causes behind diseases such as Amyotrophic Lateral Sclerosis.

© 2016 Elsevier Inc. All rights reserved.

1. Introduction

Oxidative stress is a pathological condition which implies overproduction of Reactive Oxygen Species (ROS) under conditions in which the cell's antioxidant defense system becomes no longer effective (Heather and Teismann, 2009; Sims-Robinson et al., 2013). ROS are generated as a consequence of an aerobic metabolism determined by mitochondrial respiration (Hool, 2006). To eliminate these ROS, cells develop several protective

mechanisms. Among them, enzymes such as superoxide dismutases, catalases and glutathione peroxidases directly transform some ROS into compounds of lower toxicity through the oxidation of antioxidant metabolites: reduced glutathione, thioredoxin, and ascorbic acid (Reynolds et al., 2007; Sha et al., 2013). Low level of ROS participate in cell division and growth regulation, apoptosis regulation, oxidative modifications of biomolecules in extracellular space, protection from pathogen invasion, etc. (Han et al., 2008). High levels of ROS may appear in other situations (such as mitochondrial dysfunction, excitotoxic insult, or inflammation), and cause DNA mutations, ion channel damage, intensification of the lipid peroxidation process, and oxidation of proteins and other biomolecules (Lambert et al., 2004; Ljubisavljevic, 2014; Pitt et al.,

* Corresponding author at: Departamento de Fisiología, Facultad de Farmacia, Universidad de Sevilla, C/Profesor García González no. 2, 41012 Sevilla, Spain.

E-mail address: pnunez@us.es (P. Nunez-Abades).

2000) which lead to impairing cell function. They may also affect different transcription factors, growth factors, kinases phosphatases and cytokines (Arrigo, 1999; Emerit et al., 2004; Tirosch et al., 2000; Valencia and Moran, 2004; Vimard et al., 2011). At the level of the membrane, ROS alter ATP-sensitive K^+ currents, L-type Ca^{2+} currents (Goldhaber and Liu, 1994; Racay et al., 1997) and delayed rectifier K^+ currents (Goldhaber et al., 1989), ions transporters (Kourie, 1998), either through direct oxidation of lipids, or through alterations of cell membrane proteins and intracellular signaling pathways (Hool, 2006; Zhu et al., 2005). ROS can also affect membrane properties including variations in cell cytoskeletal architecture and membrane stiffness, membrane potential, ionic gradients, action potential duration and amplitude, spontaneous activity, and excitability (Jovanovic and Jovanovic, 2013; Nakaya et al., 1992; Nani et al., 2010; Pardillo-Díaz et al., 2015; Sinha et al., 2015).

Lipid peroxidation (LPO) is one of the most commonly studied processes of redox cell signalization disorders where free radicals have a great importance (Ayala et al., 2014). Through different mechanisms, LPO disrupts membrane barrier function, inactivates membrane enzymes, and increases permeability for water, monovalent and divalent ions, and even high molecular weight compounds (Ferretti and Bacchetti, 2011; Ljubisavljevic, 2014; Nam, 2011). Cumene hydroperoxide (CH) is a stable organic oxidizing agent that is known to penetrate into the inner hydrophobic part of the membrane lipid bilayer, causing extensive peroxidation of lipids (to a much greater extent than hydrogen peroxide) (Jovanovic and Jovanovic, 2013; van den Berg et al., 1992). CH has also been described to be able to react with aminoacids and proteins with multiple effects, such as oxidation of side-chains, backbone fragmentation, dimerization/aggregation, unfolding or conformational changes, enzymatic inactivation, and alterations in cellular handling and turnover of proteins, as singlet oxygen does (Ayala et al., 2014; Davies, 2003; Gracanin et al., 2009). Consequently, CH has been used to inflict oxidative stress in vitro in neurons (Jovanovic and Jovanovic, 2013; Nakaya et al., 1992; Nani et al., 2010; Pardillo-Díaz et al., 2015). Regarding neuronal cell damage, it is widely believed that oxidative stress has a fundamental role in neuronal degeneration and might be one factor in the development of different diseases, such as Amyotrophic Lateral Sclerosis (ALS), Parkinson's, Schizophrenia, Alzheimer's (Andersen, 2004; Cabungcal et al., 2014; Cleveland and Rothstein, 2001; Reynolds et al., 2007) and ageing (Muller et al., 2007). ALS is a progressive neurodegenerative disease that results from the death of the upper motor neurons of the motor cortex, including layer pyramidal neurons, that regulate voluntary control of motor output (Mochizuki et al., 2011). In vitro and in vivo clinical and preclinical studies show that ALS is characterized by higher levels of oxidative stress biomarkers and by lower levels of antioxidant defense biomarkers in the brain and peripheral tissues (reviewed in Niedzielska et al., 2015). Post-mortem studies on tissue samples from ALS patients support the hypothesis of the oxidative damage in proteins, lipids, and DNA (Bogdanov et al., 2000; Ihara et al., 2005; Smith et al., 1998; Tohgi et al., 1999). Indeed, recent positron emission tomography (PET) imaging data in humans have confirmed that oxidative stress is enhanced in motor cortex in ALS patients compared with controls (Ikawa et al., 2014). Moreover, recent studies have demonstrated extensive early changes to the morphology of motor cortex neurons in SOD1 mice (mouse model of ALS based on oxidative stress), which thus confirms clinically relevant cortical pathophysiology more faithfully than previously thought (Fogarty et al., 2015; Saba et al., 2015). All this evidence leads to the conclusion that oxidative stress may be an important factor associated with the development of neurodegeneration in ALS patients.

In our study, we have attempted to clarify how oxidative stress evokes dose dependent changes in the functional properties of neurons. In previous studies, we have demonstrated that oxidative stress compromises both neuronal excitability and the capability of generating action potentials (Pardillo-Díaz et al., 2015). In order to understand how oxidative stress modifies neuron cell membrane properties, we have used three concentrations of CH at quantities that do not induce early cell death (Shimura et al., 1985; Vimard et al., 1996, 2011; Vroegop et al., 1995). We have registered the electrophysiological properties of pyramidal motor cortex neurons in brain slices by means of whole cell recordings. Our study aims to answer important questions such as: (i) Does acute oxidative stress induce functional changes in a dose dependent manner? (ii) In case such changes were dose dependent, are some membrane properties more sensitive to the drug concentration? (iii) Can membrane properties be recovered to resting values after washout of the drug? (iv) Can the pre-treatment with antioxidant agents, such as melatonin, prevent the LPO and the physiological changes induced by CH? (v) Are alterations in membrane properties dependent of neuronal size? The answers to these questions could be potentially relevant to explain the role oxidative damage may have in neurological disease, such as ALS.

2. Experimental procedures

The method has been described in detail in a previous publication by our lab team (Pardillo-Díaz et al., 2015). Briefly, the present study was carried out in strict accordance with the recommendations of the Guide for the Care and Use of Laboratory Animals of the European Community Directive 2003/65 and the Spanish Royal Decree 120/2005. The research protocol was approved by the Animal Ethics Committee of the University of Seville. Wistar rats (20–40 days of age) were deeply anaesthetized with chloral hydrate (4%, Panreac). Brains were quickly removed and placed in ice-cold low-calcium Artificial Cerebro-Spinal Fluid (ACSF). Transverse sections (thickness 300 μm) of the primary motor cortex were cut off on a vibratome (NVLSM1, WPI), placed in an ACSF-filled chamber for 30 min at $\sim 37^\circ\text{C}$, and then stored at $\sim 21^\circ\text{C}$. The composition of the ACSF (in mM) was as follows: 126 NaCl, 2 KCl, 1.25 Na_2HPO_4 , 26 NaHCO_3 , 10 glucose, 2 MgCl_2 , and 2 CaCl_2 . For the low-calcium-ACSF solution, the concentrations were 4 MgCl_2 mM and 0.1 CaCl_2 mM. Both ACSF and low-calcium-ACSF solutions were bubbled with 95% O_2 –5% CO_2 (pH 7.4).

2.1. Whole-cell patch-clamp recordings

Slices containing the primary motor cortex were transferred to a recording chamber and superfused at 1–2 ml min^{-1} with circulating aerated ACSF warmed to $33 \pm 1^\circ\text{C}$ via a feedback-controlled heater (TC 324B; Warner). Neurons were patch-clamped under direct visual control using a Nikon Eclipse FN1 microscope equipped with infrared-differential interference contrast (IR-DIC) optics, a 40 \times water immersion objective, and a WAT-902H2 Ultimate Camera. Cortical pyramidal neurons were distinguished by their typical morphology (Stuart et al., 1993) including a large pyramidal-shaped soma and a prominent apical dendrite extending vertically toward the pial surface. Patch pipettes were pulled (PC-10, Narishige, Tokyo, Japan) from borosilicate glass capillaries with filament (inner diameter 0.6, outer diameter 1 mm; Narishige) to a tip resistance of 3–5 $\text{M}\Omega$. Patch pipettes were filled with (in mM): 120 K-gluconate, 10 KCl, 10 phosphocreatine disodium salt, 2 MgATP, 0.3 NaGTP, 0.1 EGTA, 10HEPES, adjusted to pH 7.3 with KOH. The osmolarity of the intracellular solution was 285 mosmol/kg, adjusted with sucrose. Whole-cell recording configuration was obtained using a micro-manipulator (MP-225, Sutter) and a patch-clamp amplifier

(Multiclamp 700B, Axon Instruments, Molecular Devices, Sunnyvale, CA, USA). Giga seals ($>1\text{ G}\Omega$) were always obtained before rupture of the patch and pipette capacitance was compensated for before breaking in, and, in current-clamp mode, the bridge was periodically balanced using the auto-adjust feature. Series resistance was $20\text{ M}\Omega$ or less during recording. Current-clamp recordings were low-pass Bessel-filtered at 10 kHz ; data were digitized at $2\text{--}20\text{ kHz}$ with a Digidata 1550 analog-to-digital converter and acquired using the pCLAMP 10 software (Molecular Devices). Data were stored and subsequently prepared for analysis with the Clampfit 10.4 software (Molecular Devices).

2.2. Drugs and protocols

CH (Sigma–Aldrich, St. Louis, MO, USA) was prepared just prior to experiments from 60 mM stock solutions stored at -80°C . To avoid concerns associated with incomplete drug washout or incomplete recovery to control conditions, and, for all protocols below, only one cortical neuron per slice was recorded.

As a preliminary objective, we wanted to know if the effects demonstrated for CH at $10\ \mu\text{M}$ in our previous study (Pardillo-Díaz et al., 2015) were reversible. In that study, it was found that the application of CH ($10\ \mu\text{M}$) produces noticeable effects in some parameters, such as resistance and rheobase, at 5 min and 15 min (Pardillo-Díaz et al., 2015). For this reason, in this work, we have first tested the effects of the CH washout 5 min after administration, and we have compared the electrophysiological parameters obtained in current-clamp mode at the beginning, after 5 min CH application and 15 min after the washout. We have secondly measured the same parameters in a similar experiment in which we maintained the CH application for a longer period of 15 min duration.

A second objective of the present study was to compare the effects of CH at different concentrations: $1\ \mu\text{M}$ ($n=30$); $10\ \mu\text{M}$ ($n=30$); and $100\ \mu\text{M}$ ($n=30$). Data obtained for the $10\ \mu\text{M}$ concentration are those reported in Pardillo-Díaz et al. (2015) and were recorded in the same experimental run as the data reported here for other concentrations. Each neuron was initially superfused with normal ACSF to study their electrophysiological parameters in current-clamp mode under initial condition (time 0). After this, the slice would be superfused with ACSF containing 1, 10 or $100\ \mu\text{M}$ CH for 30 min, and voltage responses were acquired at 5, 15 and 30 min after CH application. In order to assess if long-lasting recording (30 min) affects the viability of the cells, we have also evaluated the electrophysiological properties of 20 neurons in the absence of the drug, at the same time intervals. Data from these cells were considered as a control group.

In addition, we wanted to find out the sequential effects that different concentrations of CH may have on each neuron. In order to do so, we have applied the 3 different concentrations (1, 10 and $100\ \mu\text{M}$) to the same cell at 3 moments in time. Electrophysiological parameters were analyzed before drug administration ($0\ \mu\text{M}$), and 10 min after the exposure to each of the 3 concentrations.

To determine the level of oxidative stress induced by CH and whether it can be prevented by the use of melatonin, we used a technique to measure hydroperoxide which relies upon the rapid hydroperoxide-mediated oxidation of Fe^{2+} under acidic conditions. Fe^{3+} forms a chromophore with xylenol orange which strongly absorbs at 560 nm (Jiang et al., 1991). To know the level of LPO induced by CH, trimmed brain slices (thickness $300\ \mu\text{m}$) containing primary motor cortex were placed in artificial ACSF with 1, 10 and $100\ \mu\text{M}$ of CH and they were each incubated for 5, 15 and 30 min. Slices were placed in normal ACSF to stop the chain-reaction evoked by CH and stored in HEPEs solution. The slices were homogenized during 10 s using a sonicator (Hielscher UP 100H). Then, samples were centrifuged at $2400\times g$ and the

supernatants were collected. Total protein was determined by the Lowry method. The protocol for LPO measurements was adapted for a microplate reader (Asys UVM 340). $40\ \mu\text{g}$ of protein was incubated with $90\ \mu\text{l}$ of H_2SO_4 for 30 min. After addition of $100\ \mu\text{l}$ FOX reagent (0.5 mM ferrous ammonium sulfate, 0.2 mM xylenol orange and 200 mM sorbitol in 25 mM H_2SO_4) the mixture was incubated at room temperature for 45 min, protected from light. The formation of ferric ions was detected by measuring the resulting colored complex with xylenol orange. To study if the oxidative stress can be prevented by melatonin the same protocol was used but brain slices were placed in artificial ACSF with $10\ \mu\text{M}$ of CH and $50\ \mu\text{M}$ of melatonin and they were each incubated for 5, 15 and 30 min.

Finally, to investigate if melatonin prevents the physiological changes induced by CH at $10\ \mu\text{M}$, pyramidal neurons ($n=6$) were initially superfused with normal ACSF with melatonin ($50\ \mu\text{M}$) to measure electrophysiological parameters in current-clamp mode for the initial condition (time 0). After this, each slice was superfused with ACSF with melatonin ($50\ \mu\text{M}$) and $10\ \mu\text{M}$ CH for 30 min, and voltage responses were recorded during this period.

2.3. Current-clamp recordings and analysis

All pyramidal neurons included in the analysis showed a stable resting membrane potential of -60 mV or less, an action potential higher than 80 mV , and repetitive firing in response to depolarization pulses of 1 s. The following parameters were recorded and quantified: (1) cell capacitance. (2) Resting membrane potentials were measured as the difference between the intracellular and extracellular potentials after withdrawing the recording electrode from the cell. (3) Input resistance was determined by passing positive and negative square current steps (500 ms , 1 Hz ; with 10 pA increments) and calculated as the slope of the current-voltage plot. When there was evidence of an inward rectification, or sag, the voltage value used for this latter plot was the value achieved at the peak (for details see Carrascal et al., 2005, 2006). (4) The rheobase was the minimum current injected (square pulse of 100 ms , 1 Hz ; 5 pA increments) that generated an action potential in 50% of the cases. (5) Voltage depolarization was the increase in membrane potential required for the cell to reach the spike threshold. To determine spike threshold, the action potential recording was differentiated, with the spike onset taken as the value of the membrane potential at which the first derivative exceeded 10 V s^{-1} . Voltage threshold was calculated by adding depolarization voltage to the resting membrane potential. (6) Spikes were averaged (six sweeps) to measure their characteristics: amplitude and duration ($100\ \mu\text{s}$, $5\text{--}20\text{ nA}$). Amplitude was the voltage increment between the resting level and the spike voltage peak. The value of the duration of the action potential was determined as the width of the spike at its half amplitude (see also Nunez-Abades et al., 1993). (7) The tonic component of firing was measured from the repetitive discharge evoked by depolarizing square current steps (1 s , 0.5 Hz ; $10\text{--}50\text{ pA}$ increments). The steady-state firing frequency was taken as the average number of spikes during the last 500 ms of the repetitive discharge. The relationship between the steady-state firing frequency and injected current was plotted ($f\text{--}I$ plot) and the slope (*gain*) calculated. Maximum frequency was considered as the highest frequency achieved by the neuron regardless of the current intensity, and the cancellation current was the intensity current in which the neuron would cease to fire.

2.4. Statistical analysis

Results were expressed as mean \pm standard error of the mean; n refers to the number of cells. For statistical calculations, we have

used SPSS 22.0 (IBM software, Armonk, NY, USA). Data distribution was first processed with a normality test (Shapiro–Wilk test). A paired *t*-test was used to check that the mean values (0 min and 30 min) were the same when checking cell viability under whole-cell recording configuration. An analysis of variance (ANOVA) was used to compare the means of the electrophysiological parameters for the experiments carried out to check the drug washout effects. Two-way ANOVA was used to analyze the means of the time variable (0, 5, 15 and 30 min) and for the concentration variable (0 μ M, 1 μ M, 10 μ M, and 100 μ M) for each electrophysiological parameter. ANOVA was again applied to analyze differences of the sequential application of the drug. If there were significant differences, then we used the Bonferroni test to perform pairwise comparisons between groups. Two-way ANOVA was also used to analyze the mean values of LPO at 0, 5, 15 and 30 min at the three concentrations previously used in this study (1 μ M, 10 μ M, and 100 μ M). One factor ANOVA was used to determine if there were significant differences between the group treated with CH 10 μ M and the group pre-treated with 50 μ M melatonin plus 10 μ M CH at 0, 5, 15 and 30 min. If there were significant differences, Bonferroni correction was used again to perform a comparison between

groups. Two groups of data were considered statistically different if $P \leq 0.05$. The correlation between variables was measured by the Pearson's correlation coefficient (*r*).

3. Results

We have first evaluated the electrophysiological properties of 20 neurons in order to assess whether they had been affected by a whole-cell recording configuration over sessions of at least 30 min duration. As seen in Fig. 1A, and for a representative neuron, membrane potential remained stable at approximately -74 mV throughout the recording session and the cell responded to a current injection (-100 pA) with similar changes in voltage. Membrane resistance was calculated from these voltage changes and resulted in 210, 205, 209 and 210 M Ω , for 1, 5, 15 and 30 min of the recording, respectively (see arrows in Fig. 1A). For all neurons, mean values obtained at the beginning of the recording session and at 5, 15 and 30 min did not show any significant statistical difference on membrane potential, input resistance, amplitude of the action potential, gain or maximum discharge as seen in Figs. 1, 2 and 4.




Fig. 1. Cumene hydroperoxide (CH) washout effects on membrane potential of pyramidal neurons from the motor cortex.

(A) Electrophysiological recording showing that membrane potential and voltage response to negative current pulses of -100 pA remains stable for 30 min in the control condition. (B) CH washout 5 min after drug administration (10 μ M). (C) CH washout 15 min after drug administration (10 μ M).




Fig. 2. Dose, time and size dependent effects of cumene hydroperoxide (CH) of pyramidal neurons from the motor cortex.

(A–D) Plots showing dose and time dependent effects of CH on membrane potential (A), input resistance (B), rheobase (C) and voltage threshold (D). These plots illustrate the mean values \pm standard error at 0, 5, 15 and 30 min in control condition (green; $n = 20$) and after the exposure to CH 1 μ M (blue; $n = 30$), 10 μ M (red; $n = 30$) and 100 μ M (black; $n = 30$). The asterisk indicates statistical significance in relation to the control situation and a cross represents statistical significance between correlative measures. (E and F) Relationship between capacitance and input resistance change in absolute (E) and in relative (F) values after 30 min to CH 100 μ M. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

3.1. Cumene hydroperoxide washout effects on resting membrane potential and resistance of pyramidal neurons from the motor cortex

For a representative neuron, Fig. 1B illustrates the effects of the application of 10 μ M CH on membrane potential during 5 min and then, during the washout of the drug. The recording shows how depolarization starts at 3 min reaching its highest value, of approximately 6 mV, at 8 min. Depolarization remained steady until 12 min. Then, membrane potential started a progressive repolarization, reaching to a value similar to the one recorded at

rest (-68 mV) at 15 min (10 min of the washout). Membrane potential kept that value until the end of the experiment (30 min). Repolarization during washout, as described in Fig. 1B, affected a total of 5 cells. For the other 5 cells (total population $n = 10$), only partial repolarization was observed ($<60\%$ of the change).

One of our objectives was to describe CH washout effects on membrane resistance. To accomplish this objective, we applied negative current pulses (-100 pA, 200 ms) at one-minute intervals and recorded voltage response. CH produced a biphasic change on membrane resistance, characterized by a transient increase at

5 min of drug application which then progressively fell under control values. In Fig. 1B, cell resistance shifted from 184 M Ω , in control, to 223 M Ω at 5 min. Later, it progressively diminished until reaching 180 M Ω after recording for 15 min (10 min of washout, indicated by the third arrow in the figure). The same value remained for a 20 min recording (15 min of washout). At 30 min of the recording, the value registered was 181 M Ω (fourth arrow in the figure). A total of 6/10 cells presented the same result, while the rest showed a membrane resistance with values under the ones registered at the beginning of the experiment. This could be due to a late effect of CH on resistance, and not necessarily to the washout itself.

We also looked into the effect of the washout on the firing repetitive properties. After a CH exposure of 5 min, all the cells ($n = 10$) in the study presented a decrease in gain ($55.2 \pm 12.6 \text{ AP s}^{-1} \text{ nA}^{-1}$ vs $36.7 \pm 15.5 \text{ AP s}^{-1} \text{ nA}^{-1}$). After the washout, 4 cells showed an improvement in the gain and maximum firing frequency (<40% of the change). Four neurons did not present any kind of reversibility to the effects of CH and retained a low gain and low maximum frequency. The remaining 2 cells completely lost their ability to repetitively discharge action potentials during the washout. From these experiments, we can conclude that a short exposure to the drug (5 min) allows for a partial recovery

from the effects that CH may have over membrane potential, resistance and repetitive discharge properties.

Fig. 1C shows, for a representative neuron, how the effect of 15 min CH application is the same to the one described in Fig. 1B. However, after the washout, the cell did not present any recovery regarding the parameters analyzed in the study. For the neuron depicted, membrane potential depolarized from -73 mV up to -68 mV after 15 min of CH application, and that value remained the same until the end of the recording (15 min of washout). The same applies for the entire population of neurons. The cell depicted in the figure showed a resistance of 166 M Ω in control, and then augmented to 202 M Ω after 5 min CH application. There was a decrease to 141 M Ω after 15 min. Drug washout did not produce any recovery to the initial values in resistance, on the contrary, resistance continued its decrease up to 112 M Ω at the end of the recording session. This phenomenon was observed for the whole population, with the result of a lower resistance value after the washout which is characteristic of the CH application itself. In terms of the gain and the maximum frequency, only 3 cells partially reverted to original values (30% recovery) during washout, while the remaining 7 retained a decrease in both parameters. Actually, 3 of the latter completely lost their capability of repetitively discharge and did not get recovered during washout.




Fig. 3. Effects of sequential application of increasing doses of cumene hydroperoxide (CH) on membrane resistance of pyramidal neurons from the motor cortex. (A–D) Recordings showing, in the same cell, membrane potential responses to current steps of 10 pA without CH administration (A), after 10 min of 1 μM CH administration (B), after 10 more minutes of 10 μM CH administration (C) and after 10 more minutes of 100 μM CH application (D). The black arrow in A indicates the presence of sag. (E) Relation between current and voltage response for the cell represented in A–D. (F) Histogram illustrating the mean values of membrane resistance in control condition and the other three situations for the whole population ($n = 10$).

Therefore, continuous exposure to CH (15 min) produced alterations to membrane properties that were hardly reversible.

3.2. Dose dependent effects of cumene hydroperoxide on membrane properties of pyramidal neurons from the motor cortex

Dose response effects of CH were studied during 30 min on membrane properties, namely membrane potential, input resistance, rheobase and threshold voltage. Fig. 2A shows the effects on membrane potential for control and the 3 concentrations (1, 10 and 100 μM). In the cases of control and 1 μM CH, no significant changes were observed for any of the cells along the 30 min duration of the study. However, for the case of the 10 μM CH concentration, membrane potential significantly depolarized at

5 min, and even though it reached values slightly more depolarized at 15 min and 30 min, those differences were not statistically significant when compared with the ones obtained at 5 min. When the concentration of 100 μM CH was applied, depolarization was significant at 30 min (-68.6 ± 1.3 in initial condition vs -61.9 ± 1.4 at 30 min).

Fig. 2B shows the effects on membrane resistance in a pool of neurons in control condition ($n=20$) and each one of the other 3 populations of neurons were exposed to one of the 3 drug concentrations of the study ($n=30$ for each one). As seen in the figure, membrane resistance does not change in control condition over time neither when 1 μM CH is applied during the 30 min the experiment lasted. However, input resistance showed a biphasic response for the other 2 concentrations, with a significant increase




Fig. 4. Dose and time dependent effects of cumene hydroperoxide (CH) on action potential amplitude (A), action potential duration (B), firing repetitive properties (C), gain (D), maximum frequency (E) and cancelation current (F) of pyramidal neurons from the motor cortex.

These plots illustrate the mean values \pm standard error at 0, 5, 15 and 30 min in control condition (green; $n=20$) and after the exposure to CH 1 μM (blue; $n=30$), 10 μM (red; $n=30$) and 100 μM (black; $n=30$). The asterisk indicates statistical significance in relation to the control situation and a cross represents statistical significance between correlative measures. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

at 5 min. Subsequently, we observed a significant decrease at 15 and 30 min, respectively, when compared with the initial value. More specifically, input resistance augmented at 5 min for the 100 μM CH concentration ($162.1 \pm 9.5 \text{ M}\Omega$ vs $189.2 \pm 12.4 \text{ M}\Omega$). Later, it diminished at 15 and 30 min ($110.6 \pm 9.8 \text{ M}\Omega$ and $99.7 \pm 11.1 \text{ M}\Omega$, respectively). Values at 30 min were slightly lower than the ones obtained at 15 min, but there was not a significant difference between them. The 10 and the 100 μM CH concentrations had bigger (and significant) effects than 1 μM CH at 5 min, but no differences were found between 10 and 100 μM CH.

To confirm that the 10 and 100 μM concentrations produced similar effects on resistance another experiment was carried out ($n = 10$), in which we analyzed the voltage responses to current

injections while the neuron was being sequentially exposed to increasing drug concentrations for 10 min: 0, 1, 10 and 100 μM of CH. Fig. 3 shows, for a representative neuron, that voltage responses were similar before drug application and for the 1 μM CH concentration, both demonstrating that membrane resistance was not modified in this situation. However, in the absence of the drug, the neuron (Fig. 3A) presents a voltage rectification of the membrane potential (known as “sag”) approximately at 80 ms, which disappears after 1 μM CH application (Fig. 3B). The resulting voltage responses obtained when the application was of 10 and 100 μM CH were smaller than those obtained in control, and were also accompanied by a membrane depolarization from -75 mV to -66 and -65 mV , respectively. The




Fig. 5. Effects of sequential application of increasing doses of cumene hydroperoxide (CH) on firing repetitive properties of pyramidal neurons from the motor cortex. (A–D) Recordings showing, in the same cell, voltage response to depolarizing current pulses of 1 s duration and 500 pA intensity without CH administration (A), after 10 min of 1 μM CH administration (B), after 10 more minutes of 10 μM CH administration (C) and after 10 more minutes of 100 μM CH application (D). (E) Relation between current and firing frequency for the cell represented in A–D. (F) Histogram illustrating the mean values of gain in control condition and the other three situations for the population that kept their firing properties during 30 min ($n = 6$).

measure of the membrane resistance for each neuron, before drug application and for the 3 drug concentrations, was calculated as the slope that represents the relation between current intensity and voltage response as depicted in Fig. 3A–D. For the neuron described in Fig. 3A–D, resistance was 103 M Ω without CH, 102 M Ω in 1 μ M CH. It dropped to 51 M Ω in 10 μ M CH, and to 50 M Ω in 100 μ M CH, respectively (Fig. 3E). For the whole population ($n = 10$) resistance was 170.2 \pm 6.1 M Ω in control, 165.3 \pm 6.7 M Ω in 1 μ M CH. It went down again to 135.7 \pm 5.7 M Ω in 10 μ M CH, and 120.6 \pm 11.8 M Ω in 100 μ M CH, respectively (Fig. 3F). From this data, we conclude that neurons do not evidence any alteration to membrane potential and resistance in the 1 μ M CH concentration in respect to initial values, and that the effects of the 100 μ M CH concentration do not vary significantly from the changes produced by the administration of 10 μ M CH.

3.3. Dose dependent effects of cumene hydroperoxide on rheobase and voltage threshold of pyramidal neurons from the motor cortex

The administration of CH triggered up some changes on the rheobase of pyramidal neurons from the motor cortex. Fig. 2C illustrates the effects on the rheobase which depend on the time and the dose of the CH concentration. No significant change could be found on rheobase in control condition and when 1 μ M CH was applied during 30 min. However, when the concentration applied was 10 or 100 μ M CH, the rheobase decreased significantly after 5 min application in comparison with initial condition (for example, for 100 μ M the decrement was 193.7 \pm 10.6 pA vs 149.6 \pm 15.7 pA) while it increased significantly if compared with initial condition after 15 and 30 min application (for 100 μ M, the increase was 249.9 \pm 41.6 pA and 263.1 \pm 35.3 pA). Values at 30 min were only slightly larger than those obtained at 15 min, and they did not differ significantly. Changes in rheobase were parallel to changes in resistance and, therefore, the voltage threshold remained unaltered during the 30 min duration for the 3 concentrations (Fig. 2D). As a conclusion, Ohm's Law is observable in the membrane behavior of pyramidal neurons, even under the effect of different concentrations of the drug.

Later on, we analyzed whether the effects of oxidative stress produced by CH on membrane properties would depend on neuronal size: that is to say, whether larger neurons presented a bigger sensitivity to CH or conversely. An estimation of the neuronal size can be inferred from membrane capacitance. Fig. 2E and F shows the relation between capacitance and resistance changes of neurons recorded at 100 μ M CH, in absolute values (Fig. 2E) and in relative values (Fig. 2F). Data shows a linear relation between capacitance and input resistance change when calculated from absolute values ($P < 0.001$; $r > 0.69$), and also from relative values ($P < 0.001$; $r > 0.71$). Similar results were obtained when 10 μ M CH was administered (not shown). From these figures, we can conclude that pyramidal neurons from the motor cortex that have a larger membrane surface are more likely to suffer alterations in their electrical membrane properties by oxidative stress mediated by CH.

3.4. Dose dependent effects of cumene hydroperoxide on amplitude and duration of the action potential of pyramidal neurons from the motor cortex

Fig. 4A describes the effects of CH on action potential amplitude in control condition and at the 3 different concentrations. No differences were found in control condition for this parameter while the 1 μ M CH concentration produced a gradual decrease on action potential amplitude, which became statistically significant at 30 min. For the 10 and 100 μ M concentrations, the mean decrease was more evident, reaching statistical significance at

5 min as compared to the initial condition, and at 30 min in comparison with 15 min. For the 100 μ M concentration under initial condition, the mean amplitude was 120.1 \pm 2.0, going down to 114.8 \pm 5.7 mV at 5 min, and even lower to 110.3 \pm 3.8 mV (15 min) and 91.2 \pm 3.3 mV (30 min). The highest difference between initial and final condition for the measure of the amplitude could be seen when the concentration applied was 100 μ M. More specifically, for all concentrations, amplitude differences were the following: 8.4 \pm 3.9, 18.8 \pm 2.3 and 28.9 \pm 5.8 mV, for 1, 10 and 100 μ M CH respectively. Differences of the effects on the action potential were significant between the 3 concentrations at 30 min.

In Fig. 4B, we show the effects on the duration of the action potential in control condition and for the 3 concentrations. This parameter became progressively longer after the 10 and 100 μ M CH exposure while remaining unaltered in control condition and after the 1 μ M CH application. However, statistical significance was already observable at 5 min in comparison with control condition, and 30 min in comparison with 15 min condition, for both 10 μ M and 100 μ M CH. For example, and for the administration of 100 μ M CH, the duration of the action potential was 1.50 \pm 0.11 ms (0 min), 1.82 \pm 0.13 ms (5 min), 1.87 \pm 0.28 ms (15 min), and, finally, 2.31 \pm 0.15 ms (30 min).

3.5. Dose dependent effects of cumene hydroperoxide on firing properties of pyramidal neurons from the motor cortex

The pyramidal neurons in the study exhibited a phasic-tonic discharge as a response to sustained current depolarizing pulses, not only in control, but also when the concentration applied was 1 μ M CH (see Fig. 5A and B). Nevertheless, after the administration of 10 μ M CH, 20% of the cells (15 min) and 43% (30 min) lost their ability to discharge action potentials in a repetitive way. This effect was even greater when the concentration was 100 μ M CH affecting 40% (5 min), 80% (15 min), and 100% (30 min) of the cells (Fig. 4C).

Fig. 5A–D shows, for a representative neuron ($n = 10$), the results of another experiment in which we analyzed the repetitive firing properties observed while the neuron was being sequentially exposed to each of 3 different drug concentrations for 10 min. Before drug application (0 μ M), and for the 1 and 10 μ M CH concentrations, the neuron illustrated in Fig. 5 could discharge along the whole pulse duration (1 s). However, this ability is lost after 10 min of being exposed to 100 μ M CH, as being depicted in Fig. 5D. In this figure, we can observe how the cell ceases to discharge before the pulse is over. Also in Fig. 5A–C, it can be observed how the firing frequency decreases as the concentration increases: 23 AP s⁻¹ in control, 19 AP s⁻¹ for 1 μ M CH, and 12 AP s⁻¹ for 10 μ M CH. Therefore, this data may suggest that an incremental dose alters the gain of the discharge. Fig. 5A–D also shows a decrease of amplitude of the action potential in the train, which becomes more evident for the 100 μ M CH concentration.

We then measured the gain, maximum frequency and cancellation current in those cells that retained their repetitive firing properties. Fig. 5E represents the relationship between the intensity of the current injected and the firing frequency for the cell depicted in Fig. 5A–D. In this figure, it can be observed how that relationship is linear everytime low current intensities were applied. However, when intensity increases, accommodation symptoms are appreciated. Fig. 5E shows how the neuron starts to accommodate at values close to 500 pA for 0 μ M CH, 450 pA for 1 μ M CH, and 400 pA for 10 μ M CH. This data indicates that the working intensity range decreases as the drug concentration increases. Besides, when we calculated the gain for the neuron shown in Fig. 5A–E, we could see how this parameter decreased after CH application: 53 AP s⁻¹ nA⁻¹ (0 μ M CH), 46 AP s⁻¹ nA⁻¹ (1 μ M CH), and 26 AP s⁻¹ nA⁻¹ (10 μ M CH). This result is applicable

to the rest of the population. Fig. 5F shows the mean values of the gain in the different accumulating situations (time and drug concentration): $40.1 \pm 5.9 \text{ AP s}^{-1} \text{ nA}^{-1}$ ($0 \mu\text{M}$ CH), $38.3 \pm 4.9 \text{ AP s}^{-1} \text{ nA}^{-1}$ ($1 \mu\text{M}$ CH), $27.7 \pm 8.6 \text{ AP s}^{-1} \text{ nA}^{-1}$ ($10 \mu\text{M}$ CH) and $15.3 \pm 4.1 \text{ AP s}^{-1} \text{ nA}^{-1}$ ($100 \mu\text{M}$ CH). This decrease was statistically significant for the $10 \mu\text{M}$ CH and the $100 \mu\text{M}$ CH concentrations (see Fig. 5F) when compared with initial values and between themselves.

In order to carry out a more detailed analysis of the response effects on the gain and maximum frequency, a pool of 30 neurons were exposed to one of the 3 drug concentrations tested in our study. Fig. 4D illustrates how the gain remained unchanged in control condition but progressively decreased after the administration of $1 \mu\text{M}$ CH, with statistical significance at 30 min. This decrease in gain was already significant at 15 min for the $10 \mu\text{M}$ CH, and even earlier (5 min) for the $100 \mu\text{M}$ CH concentration. For the highest concentration, the decrease in gain was significant again at 15 min when compared with 5 min. By 30 min, all cells had already lost their ability to discharge repetitively. Maximum firing frequency progressively decreased with time, as seen in Fig. 4E, but did not reach significance when $1 \mu\text{M}$ CH concentration was administered. Significance was achieved (5 min) for the 10 and the $100 \mu\text{M}$ CH concentrations. Thus, after 5 min of $100 \mu\text{M}$ CH exposure, the maximum firing frequency decreased from $27.2 \pm 3.0 \text{ AP s}^{-1}$ to $17.6 \pm 3.2 \text{ AP s}^{-1}$. The decrease in the maximum frequency was significantly lower again at 15 min and at 30 min, but only after administering $100 \mu\text{M}$ of CH.

Further evidence of the narrowing of the working intensity range, after the administration of CH, was observed when analyzing the cancelation current (Fig. 4F). This figure illustrates how the cancelation of repetitive discharge occurs at lower intensities at doses of 10 and $100 \mu\text{M}$ CH, but is unaffected in control condition and at $1 \mu\text{M}$ CH. In experiments in which the $100 \mu\text{M}$ CH concentration was tested, the average value for the population in the initial condition was $650.3 \pm 48.4 \text{ pA}$. Intensity values fell progressively to $599.3 \pm 56.5 \text{ pA}$ at 5 min, and $503.7 \pm 49.9 \text{ pA}$ at 15 min, at which level statistical significance was reached. After that, it dropped to a value of zero with the complete loss of the ability to discharge repetitively for all neurons at 30 min. Once again, the 10 and $100 \mu\text{M}$ CH concentrations had larger and more significant effects than the $1 \mu\text{M}$ and among themselves with respect to repetitive firing properties (gain, maximum frequency and cancelation current) at 30 min.

3.6. Prevention of lipid peroxidation and membrane excitability changes caused by cumene hydroperoxide by melatonin

Fig. 6A illustrates LPO expressed in percentage as the increase when compared with the control situation, as a result of the exposure of the brain slices to 0 (control), 1 , 10 and $100 \mu\text{M}$ of CH during 0 , 5 , 15 and 30 min. As seen in Fig. 6A, CH evokes LPO in a dose and time dependent manner. Longer time or higher dose caused a progressive increase in LPO, significant at 5 min at the three concentrations of CH tested in respect to the control.




Fig. 6. Prevention by melatonin of lipid peroxidation (LPO) induced by cumene hydroperoxide (CH) on pyramidal neurons from the motor cortex. (A) Graphic showing LPO values, expressed in %, at 0, 5, 15 and 30 min after the exposure of CH $1 \mu\text{M}$ (blue; $n = 6$), $10 \mu\text{M}$ (red; $n = 6$) and $100 \mu\text{M}$ (black; $n = 6$). (B) Plot showing LPO values, expressed in %, at 0, 5, 15 and 30 min after the exposure of CH $10 \mu\text{M}$ ($n = 6$) and CH $10 \mu\text{M}$ plus melatonin $50 \mu\text{M}$ ($n = 6$). (C) Electrophysiological recording showing that membrane potential and voltage response to negative current pulses of -100 pA and 500 ms remain stable along 30 min exposition to CH $10 \mu\text{M}$ + $50 \mu\text{M}$ melatonin. Arrows indicate voltage response at 5, 15 and 30 min. The letter "a" indicates statistical significance in relation to control situation, "b" represents statistical significance between a concentration and the precedent one for the same period of time, and "c" shows statistical significance between $10 \mu\text{M}$ CH application and $10 \mu\text{M}$ CH + $50 \mu\text{M}$ melatonin for the same period of time. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

However, the increased lipid peroxide level plateaued in 15 and 30 min. Furthermore, it is also observable in the fact that the level of LPO with CH 1 μM in 30 min is similar to the average value found after 1 μM CH exposure in 5 min. Although the increment in LPO is dose dependent, mainly at shorter times, those increments are not proportional to the concentration, i.e., increments in LPO found with 100 μM CH are bigger but not statistically different than those found with 10 μM CH, but they are statistically different when compared 10 vs 1 μM CH.

No effect was observed in LPO in cells treated with ACSF with 50 μM melatonin when compared with cells treated with ACSF alone. As seen in Fig. 6B, melatonin prevents LPO induced by 10 μM CH. No significant differences in LPO were found between tissues treated with melatonin alone (control of melatonin) and tissues treated with melatonin and 10 μM CH at 5, 15 or at 30 min although a moderate increase of LPO was found at 30 min. Irrespective of the time period applied (5, 15 and 30 min) the differences found between tissues treated with ACSF containing 10 μM of CH and tissues treated with ACSF containing 10 μM of CH and melatonin 50 μM , were always significant.

As melatonin prevents the LPO induced by CH, we wondered if pre-treatment with this antioxidant could act as a neuroprotector and could prevent the membrane changes induced by 10 μM CH. Fig. 6C illustrates the effects of 50 μM melatonin and 10 μM CH application on membrane potential during 30 min. As seen in this figure, no changes on membrane potential or input resistance were produced for the whole period. The same response was found in 4 of the 6 neurons studied and no changes in membrane potential, input resistance or firing properties were found. However, 2 of the 6 cells showed gain decrease, around 30 min.

4. Discussion

In the present work, we have studied early changes in membrane properties of pyramidal motor cortex neurons exposed to 3 concentrations of CH (1, 10 and 100 μM). Our data indicate that CH depresses the excitability of pyramidal motor cortex neurons decreasing input resistance, amplitude of the action potential, and gain and maximum frequency of the repetitive firing discharge. These effects were time and dose dependent. Reversibility to alterations produced by CH to membrane properties is inversely proportional to exposure time: the longer the exposure, the lower the reversibility. Among all the membrane properties analyzed, the ability to repetitively discharge action potentials was the property most affected by the administration of the drug. Our data also suggest that large neurons of the motor cortex have more sensitivity to the oxidant. Finally, we have shown that pre-treatment with melatonin, an antioxidant agent, prevents the effects induced by CH on membrane properties.

The antioxidant benefits of melatonin have been extensively described in the literature (Reiter et al., 2000; Tan et al., 2007).

4.1. Methodological considerations

Oxidative stress is caused when the production of ROS exceeds the antioxidant capacity of the tissues (Heather and Teismann, 2009; Ljubisavljevic, 2014; Sims-Robinson et al., 2013). In neuronal cell damage, it is believed that this high level of ROS makes a large contribution to neuron degeneration and might be one factor in the development of different neurological diseases, such as ALS, Parkinson's and Alzheimer's (Andersen, 2004; Reynolds et al., 2007). Our research study has considered pyramidal neurons of the layer V of the primary motor cortex. Although there are several pathophysiological mechanisms that trigger motor neuron vulnerability in ALS, oxidative stress has been described to be an important factor in the case of pyramidal neurons of the motor

cortex (Cleveland and Rothstein, 2001; Kim et al., 2014). Our results demonstrate that CH is an organic oxidant agent that induces lipidic peroxidation on the cell membrane which, consequently, results in cell alterations (Ayala et al., 2014; Nakaya et al., 1992; Vimard et al., 2011). Indeed, we have demonstrated that the effects of CH on membrane properties are due to oxidative stress because melatonin prevents the electrophysiological effects of CH. In Cai et al. (2009), malondialdehyde, a typical intermediate of lipid peroxidation, gradually damaged hippocampal neurons following a 3 h exposure at concentrations of 1, 10, 100 and 1,000 μM . According to these authors, malondialdehyde may have biochemical effects on hippocampal neurons that are time and dose dependent. A similar conclusion was reached in works that studied the dose response effects of CH toxicity on cultured cells (Shimura et al., 1985; Vimard et al., 1996, 2011; Vroegop et al., 1995). Those studies proved that cell mortality rate is low (less than 5%) when CH is applied in concentrations under 1000 μM , and with administration time under 30 min. However, they also report how mortality rate reached approximately 70% when administration times are longer (60–180 min) or when drug concentrations are larger. On the basis of this evidence, we decided to use 3 CH concentrations (1, 10 and 100 μM) and 30 min administration time, which guarantee a low mortality rate, in order to study the dose and time dependent effects of CH. We have analyzed several electrophysiological parameters during drug administration with data being recorded at 5, 15 and 30 min to identify alterations to membrane properties. Prior to that, we checked that the neurons in the motor cortex keep their membrane properties stable, under our recording conditions, for a duration of 30 min, as previously demonstrated with other pools of neurons (Nani et al., 2010). Therefore, any alteration to the physiological parameters analyzed can be considered to be the effect of the drug administration to the cells and not to the recording conditions.

4.2. Alteration of membrane properties by oxidative stress

No effects on membrane potential or resistance were observed in our study in control condition and when CH was administered at 1 μM . At that concentration, it seems that neurons are able to block the oxidative effects on the physiological properties of the membrane, although a significant level of LPO was found. A reason to explain this could be that pyramidal neurons have an efficient scavenging enzyme system that reacts rapidly with CH at a 1 μM concentration, as proposed by Jovanovic and Jovanovic (2013) when attempting to explain the effects that oxidative stress has on the electrophysiological properties of Retzius neurons of the leech. The values of LPO we found at the 1 μM concentration were under 15% of increment in respect to those registered for control. These small values are similar to those found during ageing in the hypothalamus–hypophysis, (Argüelles et al., 2011). It seems that those levels of LPO are not high enough to alter membrane properties. Contrary to the results obtained at the 1 μM concentration, our exposure of cortical neurons to CH concentrations of 10 and 100 μM already produced changes in membrane potential, membrane resistance and rheobase as a result of higher levels of LPO. An unexpected finding was that the 2 concentrations caused changes of equal magnitude in these parameters. This finding may reflect that values of LPO found when 100 μM was administered were only slightly higher, but not significant, than those obtained with 10 μM . Our results suggest that oxidative stress affected the channels and/or conductances that determine passive membrane properties (Frantseva et al., 1998; Nakaya et al., 1992; Nani et al., 2010; Pardillo-Díaz et al., 2015) along a very narrow concentration range of 1–10 μM , which resembles an “all-or-nothing” response. In any case, conductances that are affected by oxidative stress would not be those involved in determining the

voltage threshold. In our study, regardless of the CH dose used, voltage threshold remained unchanged, as occurs in neurons of the hippocampus exposed to peroxide (Pellmar, 1987).

Exposure of leech neurons to CH (0.25, 1, and 1.5 μM) extended the duration of the action potentials in a dose dependent manner (Jovanovic and Jovanovic, 2013). For example, the concentration of 0.25 μM did not significantly change the action potential within 20 min, but it was significant with 1 μM . Moreover, a higher concentration (1.5 μM) of CH caused an extreme change in the shape and duration of the action potential. In our study, we observed that the action potential diminished, while its duration increased in a progressive way along the time of exposure to CH and in a concentration dependent manner. When the CH concentration was of 1 μM , the effects on the amplitude of the action potential were evident at 30 min drug administration. However, with the 10 or 100 μM CH concentrations, effects were already significant at 5 min, showing the highest alterations in amplitude with the 100 μM CH concentration. It has been proposed that oxidative stress could be affecting conductances that participate in the repolarization of the action potential and the Na^+ ionic gradients (Angelova and Muller, 2006; Jovanovic and Jovanovic, 2013; Nakaya et al., 1992; Pardillo-Díaz et al., 2015). The capability of maintaining repetitive firing of the action potentials is probably the most dramatic effect produced by CH over the electrical properties of the membrane. Our results are similar to those obtained by Jovanovic and Jovanovic (2013) in which CH (1.5 μM , 20 min) produced a complete cancelation of firing properties in leech neurons. However, they do not coincide with the study by Nani et al. (2010) in which their H_2O_2 oxidative model (1 μM , 30 min) caused an increase in the firing of the hypoglossal motoneurons of rats. This discrepancy might be due to the fact that H_2O_2 can only exert the effects achieved by CH at concentrations 10 times higher (Vimard et al., 2011). Regardless of which channels and/or conductances are underlying action potential and repetitive firing (Pardillo-Díaz et al., 2015), in our study we have demonstrated that they are dramatically affected by oxidative stress in a concentration dependent manner, being a low concentration of CH (100 μM) high enough to completely block firing discharge.

4.3. Functional considerations

We can conclude that oxidative stress induced by CH evokes important changes, in a concentration and time dependent manner, in the neuronal excitability of rat motor cortex neurons as demonstrated in modified hippocampal cells (Oh et al., 2012). Low concentration of the drug (1 μM), but only with a long exposure, already blocks the Ih current underlying sag (Robinson and Siegelbaum, 2003) and may alter conductances involved in the action potential, and firing properties, at a drug concentration which has no biochemical, physiological or morphological effects in other neuronal populations (Jovanovic and Jovanovic, 2013; Sinha et al., 2015; Vimard et al., 2011). A concentration of 10 μM of CH produced changes in membrane potential, membrane resistance and rheobase as previously demonstrated by Pardillo-Díaz et al. (2015) in a time dependent manner. Finally, 100 μM completely blocks the ability to produce repetitive discharge of action potentials with long term exposure. Brief exposure (15 min or less) also narrow the current working range because of an increase in the rheobase and a decrease of the cancelation current. Then, we can conclude that larger concentrations of CH have similar consequences as those evoked by smaller concentrations at longer exposure times.

Present data demonstrate that effects induced by a brief exposure to CH (5 min) are partially reversible, while those produced by a long-term exposure to CH (15 min) are practically irreversible. Membrane potential and resistance presented higher

reversibility to the effects of CH. Similarly to our results, H_2O_2 induced a large depolarizing shift in membrane potential that was reversible in almost half of the thalamic neurons tested (Frantseva et al., 1998). Also the change observed in input resistance could be reversed in some cases, but not in frequency depression of spontaneous postsynaptic currents induced by H_2O_2 in hypoglossal motoneurons (Nani et al., 2010). The reversibility of some of the oxidative stress-mediated effects is difficult to explain (Pellmar, 1987). Oxygen radicals produced by CH might directly attack ion channel proteins, or LPO caused by CH might indirectly inhibit ion channel functions by altering the membrane lipid milieu surrounding the channel protein (Nakaya et al., 1992). Effects reported to be reversible may be mediated by modified lipid-protein interactions rather than by irreversible protein oxidation (Frantseva et al., 1998). Then, our results may suggest that CH effects on resting membrane potential and membrane resistance are, at least in part, a consequence of the observed LPO and, for that reason, their modifications could be partially reversed with a prompt washout of the drug. This work demonstrates that CH induces LPO in pyramidal neurons and that it can be prevented by the use of melatonin. Besides, melatonin also partially or totally prevents the physiological changes that occur in cortical neurons under CH conditions. The antioxidant benefits of melatonin have been extensively described. It has been demonstrated that melatonin has direct scavenging actions against the free radicals and related products (Reiter et al., 2000; Tan et al., 2007), acts indirectly by inducing antioxidant enzymes (Reiter et al., 2000), increases the activities of the major antioxidant enzymes (Reiter, 2000) and stimulates glutathione synthesis (Barlow-Walden et al., 1995). Melatonin has also been shown to reduce the accumulation of the major products of LPO when membranes are exposed to radical-generating agents (Reiter, 2000) and to preserve, in vivo and in vitro, protein synthesis under oxidative stress condition (Argüelles et al., 2012). Then, we cannot discard in our study that other mechanisms underlying membrane changes induced by CH, apart from LPO, such as protein oxidation, can also be prevented by melatonin. However, since 50 μM melatonin has not totally prevented the effects of long time exposition to CH in some cells, additional trials with melatonin alone (a higher dose) and in combination with other drugs, such as thiol-reducing agents like N-acetylcysteine or β -mercaptoethanol (Cabungcal et al., 2014; Vimard et al., 2011), or other anti-oxidant substances, such as vitamin C (Chang et al., 2008), are needed to clarify its potential benefit. Nevertheless, we propose that neuroprotective agents should be administered before membrane properties of neurons start to change, given the low reversibility of the effects caused by oxidative stress when this is maintained in time.

In motoneurons, cell capacitance has been used as an estimation for neuronal size (Viana et al., 1994). In other words, larger neurons would present a higher capacitance. Our results provide the first experimental evidence that neurons mostly affected by oxidative stress are the largest ones, those with the highest values of capacitance. Pyramidal neurons of the motor cortex are rather large in comparison with neurons from other layers of the motor cortex and from other brain nuclei (Oswald et al., 2013). Furthermore, regarding pyramidal neurons from the motor cortex, as from other cortical areas, it has been demonstrated that morphological parameters revealed clear differences between large soma, thick shafted, large tufted corticofugal neurons and small soma, slender shafted, small tufted commissural neurons (Chagnac-Amitai et al., 1990; Mason and Larkman, 1990; Oswald et al., 2013). We propose that corticofugal neurons could be the most affected by oxidative stress in our study. If we considered oxidative stress to be the main cause of neuronal death in neurodegenerative pathologies (Andersen, 2004; Cabungcal et al., 2014; Cleveland and Rothstein, 2001; Niedzielska et al., 2015;

Parakh et al., 2013; Reynolds et al., 2007), our findings (large size and high sensitivity to oxidative stress even a low concentration) would serve to explain how pyramidal neurons of the motor cortex, in particular corticofugal neurons, also have the highest death rate levels in diseases such as ALS.

Acknowledgements

This work was partially supported by ERDF (European Regional Development Fund) UNSE10-1E-0949. Mr. Ricardo Pardillo-Díaz obtained a research fellowship funded by CACOF. We sincerely thank The Centre of Research Technology and Innovation of the University of Seville (CITIUS) for their technical support, and Ms. Rosa María Andrade García for her technical assistance. Gratitude to Professor Juan Ramón Lacalle Remigio is expressed for his advice on statistical analysis.

References

- Andersen, J.K., 2004. Oxidative stress in neurodegeneration: cause or consequence? *Nat. Med. Suppl.* 10, S18–S25.
- Angelova, P., Muller, W., 2006. Oxidative modulation of the transient potassium current IA by intracellular arachidonic acid in rat CA1 pyramidal neurons. *Eur. J. Neurosci.* 23, 2375–2384.
- Argüelles, S., Cano, M., Machado, A., Ayala, A., 2011. Effect of aging and oxidative stress on elongation factor-2 in hypothalamus and hypophysis. *Mech. Ageing Dev.* 132, 55–64.
- Argüelles, S., Muñoz, M.F., Cano, M., Machado, A., Ayala, A., 2012. In vitro and in vivo protection by melatonin against the decline of elongation factor-2 caused by lipid peroxidation: preservation of protein synthesis. *J. Pineal Res.* 53, 1–10.
- Arrigo, A.P., 1999. Gene expression and the thiol redox state. *Free Radic. Biol. Med.* 27, 936–944.
- Ayala, A., Muñoz, M.F., Argüelles, S., 2014. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid. Med. Cell Longev.* 360438.
- Bogdanov, M., Brown, R.H., Matson, W., Smart, R., Hayden, D., O'Donnell, H., Flint Beal, M., Cudkovic, M., 2000. Increased oxidative damage to DNA in ALS patients. *Free Radic. Biol. Med.* 29, 652–658.
- Cai, J., Chen, J., He, H., Yin, Z., Zhu, Z., Yin, D., 2009. Carbonyl stress: malondialdehyde induces damage on rat hippocampal neurons by disturbance of Ca²⁺ homeostasis. *Cell Biol. Toxicol.* 25, 435–445.
- Barlow-Walden, L.R., Reiter, R.J., Abe, M., Pablos, M., Menendez-Pelaez, A., Chen, L.D., Poeggeler, B., 1995. Melatonin stimulates brain glutathione peroxidase activity. *Neurochem. Int.* 26, 497–502.
- Cabungan, J.H., Counotte, D.S., Lewis, E.M., Tejada, H.A., Piantadosi, P., Pollock, C., Calhoun, G.G., Sullivan, E.M., Presgraves, E., Kil, J., Hong, L.E., Cuenod, M., Do, K.Q., O'Donnell, P., 2014. Juvenile antioxidant treatment prevents adult deficits in a developmental model of schizophrenia. *Neuron* 83, 1073–1084.
- Carrascal, L., Nieto-Gonzalez, J.L., Cameron, W.E., Torres, B., Nunez-Abades, P.A., 2005. Changes during the postnatal development in physiological and anatomical characteristics of rat motoneurons studied in vitro. *Brain Res. Brain Res. Rev.* 49, 377–387.
- Carrascal, L., Nieto-Gonzalez, J.L., Nunez-Abades, P., Torres, B., 2006. Temporal sequence of changes in electrophysiological properties of oculomotor motoneurons during postnatal development. *Neuroscience* 140, 1223–1237.
- Chagnac-Amitai, Y., Luhmann, H.J., Prince, D.A., 1990. Burst generating and regular spiking layer 5 pyramidal neurons of rat neocortex have different morphological features. *J. Comp. Neurol.* 296, 598–613.
- Chang, H.M., Huang, Y.L., Lan, C.T., Wu, U.I., Hu, M.E., Youn, S.C., 2008. Melatonin preserves superoxide dismutase activity in hypoglossal motoneurons of adult rats following peripheral nerve injury. *J. Pineal Res.* 44, 172–180.
- Cleveland, D.W., Rothstein, J.D., 2001. From Charcot to Lou Gehrig: deciphering selective motor neuron death in ALS. *Nat. Rev. Neurosci.* 2, 806–819.
- Davies, M.J., 2003. Singlet oxygen-mediated damage to proteins and its consequences. *Biochem. Biophys. Res. Commun.* 305, 761–770.
- Emerit, J., Edeas, M., Bricaire, F., 2004. Neurodegenerative diseases and oxidative stress. *Biomed. Pharmacother.* 58, 39–46.
- Ferretti, G., Bacchetti, T., 2011. Peroxidation of lipoproteins in multiple sclerosis. *J. Neurol. Sci.* 311, 92–97.
- Fogarty, M.J., Noakes, P.G., Bellingham, M.C., 2015. Motor cortex layer V pyramidal neurons exhibit dendritic regression, spine loss, and increased synaptic excitation in the presymptomatic hSOD1(G93A) mouse model of amyotrophic lateral sclerosis. *J. Neurosci.* 35, 643–647.
- Frantseva, M.V., Perez Velazquez, J.L., Carlen, P.L., 1998. Changes in membrane and synaptic properties of thalamocortical circuitry caused by hydrogen peroxide. *J. Neurophysiol.* 80, 1317–1326.
- Goldhaber, J.I., Ji, S., Lamp, S.T., Weiss, J.N., 1989. Effects of exogenous free radicals on electromechanical function and metabolism in isolated rabbit and guinea pig ventricle. Implications for ischemia and reperfusion injury. *J. Clin. Invest.* 83, 1800–1809.
- Goldhaber, J.I., Liu, E., 1994. Excitation-contraction coupling in single guinea-pig ventricular myocytes exposed to hydrogen peroxide. *J. Physiol.* 477 (Pt 1), 135–147.
- Gracani, M., Hawkins, C.L., Pattison, D.I., Davies, M.J., 2009. Singlet-oxygen-mediated amino acid and protein oxidation: formation of tryptophan peroxides and decomposition products. *Free Radic. Biol. Med.* 47, 92–102.
- Han, M.H., Hwang, S.I., Roy, D.B., Lundgren, D.H., Price, J.V., Ousman, S.S., Fernald, G.H., Gerlitz, B., Robinson, W.H., Baranzini, S.E., Grinnell, B.W., Raine, C.S., Sobel, R.A., Han, D.K., Steinman, L., 2008. Proteomic analysis of active multiple sclerosis lesions reveals therapeutic targets. *Nature* 451, 1076–1081.
- Heather, L.M., Teismann, P., 2009. Glutathione—a review on its role and significance in Parkinson's disease. *FASEB J.* 23, 3263–3272.
- Hool, L.C., 2006. Reactive oxygen species in cardiac signalling: from mitochondria to plasma membrane ion channels. *Clin. Exp. Pharmacol. Physiol.* 33, 146–151.
- Ihara, Y., Nobukuni, K., Takata, H., Hayabara, T., 2005. Oxidative stress and metal content in blood and cerebrospinal fluid of amyotrophic lateral sclerosis patients with and without a Cu, Zn superoxide dismutase mutation. *Neurol. Res.* 27, 105–108.
- Ikawa, M., Okazawa, H., Tsujikawa, T., Muramatsu, T., Kishitani, T., Kamisawa, T., Matsunaga, A., Yamamura, O., Mori, T., Hamano, T., Kiyono, Y., Nakamoto, Y., Yoneda, M., 2014. Increased cerebral oxidative stress in amyotrophic lateral sclerosis: a 62Cu-ATSM pet study. *Neurology* 84, 2033–2039.
- Jiang, Z.Y., Woollard, A.C., Wolff, S.P., 1991. Lipid hydroperoxide measurement by oxidation of Fe2p in the presence of xylenol orange: comparison with the TBA assay and an iodometric method. *Lipids* 26, 853–856.
- Jovanovic, Z., Jovanovic, S., 2013. Comparison of the effects of cumene hydroperoxide and hydrogen peroxide on Retzius nerve cells of the leech *Haemopsis sanguisuga*. *Exp. Anim.* 62, 9–17.
- Kim, C., Lee, H.C., Sung, J.J., 2014. Amyotrophic lateral sclerosis – cell based therapy and novel therapeutic development. *Exp. Neurobiol.* 23, 207–214.
- Kourie, J.I., 1998. Interaction of reactive oxygen species with ion transport mechanisms. *Am. J. Physiol.* 275, C1–24.
- Lambert, A.J., Portero-Otin, M., Pamplona, R., Merry, B.J., 2004. Effect of ageing and caloric restriction on specific markers of protein oxidative damage and membrane peroxidizability in rat liver mitochondria. *Mech. Ageing Dev.* 125, 529–538.
- Ljubisavljevic, S., 2014. Oxidative stress and neurobiology of demyelination. *Mol. Neurobiol.* doi:http://dx.doi.org/10.1007/s12035-014-9041-x.
- Mason, A., Larkman, A., 1990. Correlations between morphology and electrophysiology of pyramidal neurons in slices of rat visual cortex. II. *Electrophysiology. J. Neurosci.* 10, 1415–1428.
- Mochizuki, Y., Mizutani, T., Shimizu, T., Kawata, A., 2011. Proportional neuronal loss between the primary motor and sensory cortex in amyotrophic lateral sclerosis. *Neurosci. Lett.* 503, 73–75.
- Muller, F.L., Lustgarten, M.S., Jang, Y., Richardson, A., Van Remmen, H., 2007. Trends in oxidative aging theories. *Free Radic. Biol. Med.* 43, 477–503.
- Nakaya, H., Takeda, Y., Tohse, N., Kanno, M., 1992. Mechanism of the membrane depolarization induced by oxidative stress in guinea-pig ventricular cells. *J. Mol. Cell Cardiol.* 24, 523–534.
- Nam, T.G., 2011. Lipid peroxidation and its toxicological implications. *Toxicol. Res.* 27, 1–6.
- Nani, F., Cifra, A., Nistri, A., 2010. Transient oxidative stress evokes early changes in the functional properties of neonatal rat hypoglossal motoneurons in vitro. *Eur. J. Neurosci.* 31, 951–966.
- Niedzielska, E., Smaga, I., Gawlik, M., Moniczewski, A., Stankowicz, P., Pera, J., Filip, M., 2015. Oxidative stress in neurodegenerative diseases. *Mol. Neurobiol.* [Epub ahead of print].
- Nunez-Abades, P.A., Spielmann, J.M., Barrionuevo, G., Cameron, W.E., 1993. In vitro electrophysiology of developing genioglossal motoneurons in the rat. *J. Neurophysiol.* 70, 1401–1411.
- Oh, J.M., Choi, E.-K., Carp, R.I., Kim, Y.-S., 2012. Oxidative stress impairs autophagic flux in prion protein-deficient hippocampal cells. *Autophagy* 8 (10), 1448–1461.
- Oswald, M.J., Tantrigama, M.L., Sonntag, I., Hughes, S.M., Empson, R.M., 2013. Diversity of layer 5 projection neurons in the mouse motor cortex. *Front. Cell. Neurosci.* 7, 174.
- Parakh, S., Spencer, D.M., Halloran, M.A., Soo, K.Y., Atkin, J.D., 2013. Redox regulation in amyotrophic lateral sclerosis. *Oxid. Med. Cell Longev.* 40868, 1. doi:http://dx.doi.org/10.1155/2013/408681.
- Pardillo-Díaz, R., Carrascal, L., Ayala, A., Nunez-Abades, P., 2015. Oxidative stress induced by cumene hydroperoxide evokes changes in neuronal excitability of rat motor cortex neurons. *Neuroscience* 289, 85–98.
- Pellmar, T.C., 1987. Peroxide alters neuronal excitability in the CA1 region of guinea-pig hippocampus in vitro. *Neuroscience* 23, 447–456.
- Pitt, D., Werner, P., Raine, C.S., 2000. Glutamate excitotoxicity in a model of multiple sclerosis. *Nat. Med.* 6, 67–70.
- Racay, P., Kaplan, P., Mezesova, V., Lehotsky, J., 1997. Lipid peroxidation both inhibits Ca(2+)-ATPase and increases Ca2+ permeability of endoplasmic reticulum membrane. *Biochem. Mol. Biol. Int.* 41, 647–655.
- Reiter, R.J., 2000. Melatonin: lowering the high price of free radicals. *News Physiol. Sci.* 15, 246–250.
- Reiter, R.J., Tan, D.X., Osuna, C., Gitto, E., 2000. Actions of melatonin in the reduction of oxidative stress. A review. *J. Biomed. Sci.* 7, 444–458.
- Reynolds, A., Laurie, C., Mosley, R.L., Gendelman, H.E., 2007. Oxidative stress and the pathogenesis of neurodegenerative disorders. *Int. Rev. Neurobiol.* 82, 297–325.
- Robinson, R.B., Siegelbaum, S.A., 2003. Hyperpolarization-activated cation currents: from molecules to physiological function. *Annu. Rev. Physiol.* 65, 453–480.

- Saba, L., Viscomi, M.T., Caioli, S., Pignataro, A., Bisicchia, E., Pieri, M., Molinari, M., Ammassari-Teule, M., Zona, C., 2015. Altered functionality, morphology, and vesicular glutamate transporter expression of cortical motor neurons from a presymptomatic mouse model of Amyotrophic Lateral Sclerosis. *Cereb. Cortex* [Epub ahead of print].
- Sha, W., Martins, A.M., Laubenbacher, R., Mendes, P., Shulaev, V., 2013. The genome-wide early temporal response of *Saccharomyces cerevisiae* to oxidative stress induced by cumene hydroperoxide. *PLoS One* 8, e74939.
- Shimura, J., Shimura, E., Hosoya, N., 1985. Functional disability of rat splenocytes provoked to lipid peroxidation by cumene hydroperoxide. *Biochim. Biophys. Acta* 845, 43–47.
- Sims-Robinson, C., Hur, J., Hayes, J.M., Dauch, J.R., Keller, P.J., Brooks, S.V., Feldman, E. L., 2013. The role of oxidative stress in nervous system aging. *PLoS One* 8, e68011. doi:<http://dx.doi.org/10.1371/journal.pone.0068011>.
- Sinha, A., Chu, T.T.T., Dao, M., Chandramohanadas, R., 2015. Single-cell evaluation of red blood cell bio-mechanical and nano-structural alterations upon chemically induced oxidative stress. *Sci. Rep.* 59768 doi:<http://dx.doi.org/10.1038/srep09768>.
- Smith, R.G., Henry, Y.K., Mattson, M.P., Appel, S.H., 1998. Presence of 4-hydroxynonenal in cerebrospinal fluid of patients with sporadic amyotrophic lateral sclerosis. *Ann. Neurol.* 44, 696–699.
- Stuart, G.J., Dodt, H.U., Sakmann, B., 1993. Patch-clamp recordings from the soma and dendrites of neurons in brain slices using infrared video microscopy. *Pflugers Arch.* 423, 511–518.
- Tan, D.X., Manchester, L.C., Terron, M.P., Flores, L.J., Reiter, R.J., 2007. One molecule, many derivatives: a never-ending interaction of melatonin with reactive oxygen and nitrogen species? *J. Pineal Res.* 42, 28–42.
- Tirosh, O., Sen, C.K., Roy, S., Packer, L., 2000. Cellular and mitochondrial changes in glutamate-induced HT4 neuronal cell death. *Neuroscience* 97, 531–541.
- Tohgi, H., Abe, T., Yamizaki, K., Murata, T., Ishizaki, E., Isobe, C., 1999. Remarkable increase in cerebrospinal fluid 3-nitrotyrosine in patients with sporadic amyotrophic lateral sclerosis. *Ann. Neurol.* 46, 129–131.
- Valencia, A., Moran, J., 2004. Reactive oxygen species induce different cell death mechanisms in cultured neurons. *Free Radic. Biol. Med.* 36, 1112–1125.
- van den Berg, J.J., Op den Kamp, J.A., Lubin, B.H., Roelofsen, B., Kuypers, F.A., 1992. Kinetics and site specificity of hydroperoxide-induced oxidative damage in red blood cells. *Free Radic. Biol. Med.* 12, 487–498.
- Viana, F., Bayliss, D.A., Berger, A.J., 1994. Postnatal changes in rat hypoglossal motoneuron membrane properties. *Neuroscience* 59, 131–148.
- Vimard, F., Nouvelot, A., Duval, D., 1996. Cytotoxic effects of an oxidative stress on neuronal-like pheochromocytoma cells (PC12). *Biochem. Pharmacol.* 51, 1389–1395.
- Vimard, F., Saucet, M., Nicole, O., Feuilloley, M., Duval, D., 2011. Toxicity induced by cumene hydroperoxide in PC12 cells: protective role of thiol donors. *J. Biochem. Mol. Toxicol.* 25, 205–215.
- Vroegop, S.M., Decker, D.E., Buxser, S.E., 1995. Localization of damage induced by reactive oxygen species in cultured cells. *Free Radic. Biol. Med.* 18, 141–151.
- Zhu, D., Tan, K.S., Zhang, X., Sun, A.Y., Sun, G.Y., Lee, J.C., 2005. Hydrogen peroxide alters membrane and cytoskeleton properties and increases intercellular connections in astrocytes. *J. Cell Sci.* 118, 3695–3703.